

Ambient Air Quality Monitoring at Sunshine Canyon Landfill and Van Gogh Elementary School

Continuous monitoring of particulate matter, black carbon, wind speed, and wind direction began at the Sunshine Canyon Landfill (Landfill Site) and at Van Gogh Elementary School (Community Site) in Granada Hills in fall 2007.

These data are used to characterize ambient air pollution concentrations on a neighborhood scale in the context of the Los Angeles basin and to evaluate the impact of landfill operations on air quality in the community.

Particulate Matter (PM₁₀)

PM₁₀ is particulate matter less than 10 microns in diameter. A human hair is about 100 micrometers in diameter, and its width could hold roughly 10 PM₁₀ particles. PM₁₀ is present in dust, smoke, soot, and dirt. It can be inhaled and drawn into the lungs, causing health problems for some people.

Black Carbon (BC)

Black carbon is a sooty black material emitted from gas and diesel engines, coal-fired power plants, and other sources that burn fossil fuel. Many BC particles are too small to be visible. BC emissions can cause adverse health and climate effects.

Wind

Wind speed and wind direction are measured because they can significantly affect when and how far airborne pollutants travel from their sources.



Wind-Blown
Dust



Landfill
Operations



Dirt Roads




Vehicular
Traffic



Diesel
Engines



Industrial
Activities



What Have We Learned in Eighteen Years of Monitoring?

PM₁₀ Exceedances

Total Exceedances
over 18 Years

59

Landfill Site
Federal
Exceedances

6

Community
Site Federal
Exceedances

1,400

Landfill Site
State
Exceedances

324

Community Site
State
Exceedances

Total Exceedances in
Year 18

1

Landfill Site
Federal
Exceedances

0


Community
Site Federal
Exceedances

18

Landfill Site
State
Exceedances

4

Community Site
State
Exceedances



Air pollution from the surrounding region appears to be the main driver of high PM₁₀ concentrations at the Community Site.

High pollution levels at the Landfill Site did not occur at the same times as high pollution levels at the Community Site. Typically, when regional pollution levels were relatively low and levels at the Landfill Site were high, levels at the Community Site were not significantly high.

Landfill Site

The Landfill Site's PM₁₀ Federal and State exceedances are accompanied by high wind speeds, with wind direction falling within a narrow sector that encompasses the active portion of the landfill. While it's assumed landfill activity played a key role in federal exceedances at the Landfill Site, the absence of an upwind landfill monitoring site makes it difficult to quantify landfill contributions to neighborhood-scale pollutant concentrations.

State exceedance days at the Landfill Site are also accompanied by low-speed winds from the Los Angeles basin (south and southeast), suggesting that the addition of elevated concentrations within the basin can push the Landfill Site's PM₁₀ concentrations over the State threshold.

Community Site

On days when PM₁₀ concentrations exceed the State standard at the Community Site, wind speeds are relatively low and wind direction is predominantly from the Los Angeles basin (southeast). This suggests that regional contributions are the main driver of exceedances of the State PM₁₀ standard at the Community Site. Further, the annual average PM₁₀ concentration at the Community Site has been steadily decreasing since Year 7. This is an important finding that shows that, while PM₁₀ concentrations measured at the Landfill Site remain high relative to the regional sites over the 18-yr period, PM₁₀ concentrations measured at the Community Site are trending down.

More Key Insights in PM₁₀ and BC Monitoring at the Landfill and Community Sites

- The greatest difference in PM₁₀ and BC concentrations between the Landfill and Community sites was observed during periods of highest activity levels (i.e., typical working hours on working days).
- BC concentrations were higher at the Community Site than at the Landfill Site when the wind was from the south coast air basin in the working hour categories.
- BC concentrations at the Landfill and Community sites are lower than those measured at the four sites in the Multiple Air Toxics Exposure Study V (MATES V) during the same time period.

Eighteenth Annual Report of Ambient Air Quality Monitoring at Sunshine Canyon Landfill and Van Gogh Elementary School:

An Eighteen-Year Summary:

November 22, 2007–November 21, 2025




Annual Report Prepared for

Planning Department, City of Los Angeles
and
Los Angeles County Department of
Regional Planning
Los Angeles, California

July 2026

SPHEROS
ENVIRONMENTAL

This document contains blank pages to accommodate two-sided printing.



Eighteenth Annual Report of Ambient Air Quality Monitoring at Sunshine Canyon Landfill and Van Gogh Elementary School

An Eighteen-Year Summary November 22, 2007–November 21,
2025

Submitted by

Charles Scarborough
Bryan Penfold
Hilary Hafner
Kevin Smith

Spheros Environmental
1221 Auraria Pkwy
Denver, CO 80204-1836
Ph 303.301.7789
spherosenv.com

Prepared for

Planning Department, City of Los Angeles
City Hall, Room 525
200 N. Spring St.
Los Angeles, CA 90012
and
Los Angeles County Department of Regional
Planning
320 West Temple Street, 13th Floor
Los Angeles, CA 90012

Annual Report

SPH-1925037-10035

July 20, 2026

Contents

Figures	iv
Tables.....	vii
Acronyms.....	viii
Executive Summary.....	ix
1. Introduction.....	1-1
1.1 Baseline Year and Continuous Monitoring.....	1-1
1.2 Report Overview	1-2
1.3 Methods and Operations Background.....	1-4
2. Data Completeness.....	2-1
3. PM₁₀ Exceedances.....	3-1
3.1 Federal Exceedances	3-2
3.2 State Exceedances.....	3-9
4. Regional Comparisons of PM₁₀.....	4-1
5. PM₁₀ and BC: Effects of Wind Direction and Work Activity Levels	5-1
5.1 General Wind Roses for the Monitoring Sites.....	5-1
5.2 Wind Direction Sectors for Categorizing Data.....	5-4
5.3 Working and Non-Working Days and Hours for Categorizing Data.....	5-7
5.4 PM ₁₀ Concentrations.....	5-7
5.5 BC Concentrations.....	5-16
6. Quantitative Estimates of Landfill Impacts on Ambient Concentrations of PM₁₀ and BC.....	6-1
7. Routine Field Operations	7-1
Appendix A. Regional Concentrations of Black Carbon	A-1
Appendix B. Additional Analyses	B-1
Appendix C. Comparison of Ambient Air Toxics Concentrations to the Final Supplemental Environmental Impact Report.....	C-1

Figures

Figure 1-1. Locations of the Landfill and Community sites in relation to the three South Coast AQMD PM ₁₀ sites and four MATES V BC sites used for regional comparisons.....	1-3
Figure 3-1. The number of federal exceedances of 24-hr PM ₁₀ at the Landfill, Community, and regional monitoring sites by year over the 18-yr study period.....	3-2
Figure 3-2. PM ₁₀ concentrations at air monitoring sites across the Los Angeles area on January 7, 2025	3-3
Figure 3-3. Hourly time series of PM ₁₀ and BC concentrations (top two rows) and WS and WD measurements recorded at the Landfill and Community sites five days before and after the January 7, 2025, PM ₁₀ exceedance at the Landfill site	3-4
Figure 3-4. Wind rose from federal exceedance days during 18 continuous monitoring years at the Landfill and Community sites.....	3-8
Figure 3-5. Number of state exceedances of 24-hr PM ₁₀ at the Landfill, Community, and regional monitoring sites by year over the 18-yr study period.....	3-9
Figure 3-6. Wind rose from state 24-hr PM ₁₀ exceedance days during 18 continuous monitoring years at the Landfill and Community sites.....	3-13
Figure 4-1. Monthly average PM ₁₀ concentrations for the Landfill, Landfill North (in operation from June 2016 until May 2017), and Community sites, and three regional monitoring sites for 2008–2025	4-3
Figure 4-2. Monthly average PM ₁₀ concentrations for the Landfill and Community sites, and three regional monitoring sites from Year 13 through Year 18.....	4-3
Figure 4-3. Rolling 3-yr annual average PM ₁₀ concentrations for the Landfill and Community sites, and three regional monitoring sites for Monitoring Year 1–Year 18.....	4-4
Figure 5-1. Wind roses for the Landfill site over the 18 years of monitoring data.....	5-2
Figure 5-2. Wind roses for the Community Site over the 18 years of monitoring data	5-3
Figure 5-3. Top panel: pollution rose of hourly BC concentration at the Community site.	5-4
Figure 5-4. Aerial image of the Sunshine Canyon Landfill and the surrounding area, showing the wind direction sectors representing the landfill source used to select data for analysis from the Landfill monitor and the Community monitor	5-5
Figure 5-5. Aerial image of the Sunshine Canyon Landfill and the northern portion of the SoCAB, showing the wind direction sector representing the SoCAB source used to select data for analysis to compare with the landfill wind direction sectors depicted in Figure 5-4	5-6
Figure 5-6. Instructions for interpreting notched box-whisker plots.....	5-8
Figure 5-7. Notched box whisker plots of 18-yr hourly PM ₁₀ concentrations for northerly (“From Landfill”) wind sectors for working days, non-working days, working hours, and non-working hours within those days for the Landfill and Community monitor sites	5-10

Figure 5-8. Notched box whisker plots of 18-yr hourly PM ₁₀ concentrations for southerly wind sectors for working days, non-working days, working hours, and non-working hours within those days for the Landfill and Community monitor sites	5-11
Figure 5-9. Notched box whisker plots of Year 18 (2025) hourly PM ₁₀ concentrations for northerly wind sectors for working days, non-working days, working hours, and non-working hours within those days for the Landfill and Community monitor sites	5-12
Figure 5-10. Notched box whisker plots of Year 18 (2025) hourly PM ₁₀ concentrations for southerly wind sectors for working days, non-working days, working hours, and non-working hours within those days for the Landfill and Community monitor sites	5-13
Figure 5-11. Eighteen-yr hourly PM ₁₀ median concentrations for northerly wind sectors for working days, non-working days, working hours, and non-working hours within those days for the Landfill and Community monitor sites	5-14
Figure 5-12. Eighteen-yr hourly PM ₁₀ median concentrations for northerly wind sectors for working days, non-working days, working hours, and non-working hours within those days for the Landfill and Community monitor sites	5-14
Figure 5-13. Year 18 (2025) hourly PM ₁₀ median concentrations for northerly wind sectors for working days, non-working days, working hours, and non-working hours within those days for the Landfill and Community monitor sites	5-15
Figure 5-14. Year 18 (2025) hourly PM ₁₀ median concentrations for northerly wind sectors for working days, non-working days, working hours, and non-working hours within those days for the Landfill and Community monitor sites	5-15
Figure 5-15. Hourly PM ₁₀ median concentrations for (a) Burbank and (b) Los Angeles for working days, non-working days, working hours, and non-working hours	5-16
Figure 5-16. Notched box whisker plots of 18-yr hourly average BC concentrations for northerly wind sectors for working days, non-working days, working hours, and non-working hours within those days for the Landfill and Community monitor sites	5-18
Figure 5-17. Notched box whisker plots of 18-yr hourly average BC concentrations for southerly wind sectors for working days, non-working days, working hours, and non-working hours within those days for the Landfill and Community monitor sites	5-19
Figure 5-18. Notched box whisker plots of Year 18 (2025) hourly average BC concentrations for northerly wind sectors for working days, non-working days, working hours, and non-working hours within those days for the Landfill and Community monitor sites	5-20
Figure 5-19. Notched box whisker plots of Year 18 (2025) hourly average BC concentrations for southerly wind sectors for working days, non-working days, working hours, and non-working hours within those days for the Landfill and Community monitor sites	5-21
Figure 5-20. Eighteen-yr hourly median BC concentrations for northerly wind sectors for for working days, non-working days, working hours, and non-working hours within those days for the Landfill and Community monitor sites	5-22

Figure 5-21. Eighteen-yr hourly median BC concentrations for northerly wind sectors for working days, non-working days, working hours, and non-working hours within those days for the Landfill and Community monitor sites.....	5-22
Figure 5-22. Year 18 (2025) hourly median BC concentrations for northerly wind sectors for working days, non-working days, working hours, and non-working hours within those days for the Landfill and Community monitor sites.....	5-23
Figure 5-23. Year 18 (2025) hourly median BC concentrations for northerly wind sectors for working days, non-working days, working hours, and non-working hours within those days for the Landfill and Community monitor sites.....	5-23
Figure 6-1. Median, mean, and standard deviation of PM ₁₀ concentration differences at the Community site versus the Landfill site for northerly wind sectors for working days, non-working days, working hours, and non-working hours within those days for the Landfill and Community monitor sites based on the 18-yr monitoring period dataset.	6-3
Figure 6-2. Median, mean, and standard deviation of PM ₁₀ concentration differences at the Community site versus the Landfill site for southerly wind sectors for working days, non-working days, working hours, and non-working hours based on the 18-yr monitoring period dataset.....	6-4
Figure 6-3. Median, mean, and standard deviation of PM ₁₀ concentration differences at the Community site versus the Landfill site for northerly wind sectors for working days, non-working days, working hours, and non-working hours based on the Year 18 dataset.....	6-5
Figure 6-4. Median, mean, and standard deviation of PM ₁₀ concentration differences at the Community site versus the Landfill site for southerly wind sectors for working days, non-working days, working hours, and non-working hours based on the Year 18 (2025) dataset.....	6-6
Figure 6-5. Median, mean, and standard deviation of BC concentration differences at the Community site versus the Landfill site for northerly wind sectors for working days, non-working days, working hours, and non-working hours based on the 18-yr monitoring period dataset.....	6-7
Figure 6-6. Median, mean, and standard deviation of BC concentration differences at the Community site versus the Landfill site for southerly wind sectors for working days, non-working days, working hours, and non-working hours based on the 18-yr monitoring period dataset.....	6-8
Figure 6-7. Median, mean, and standard deviation of BC concentration differences at the Community site versus the Landfill site for northerly wind sectors for working days, non-working days, working hours, and non-working hours based on the Year 18 dataset.....	6-9
Figure 6-8. Median, mean, and standard deviation of BC concentration differences at the Community site versus the Landfill site for southerly wind sectors for working days, non-working days, working hours, and non-working hours based on the Year 18 dataset.....	6-10

Tables

Table 2-1. Data completeness statistics for hourly data during Years 1-18 of continuous monitoring and overall 18-yr averages.....	2-2
Table 3-1. Total number of 24-hr federal ($150\ \mu\text{g}/\text{m}^3$) and state ($50\ \mu\text{g}/\text{m}^3$) PM_{10} exceedances at the Landfill, Community, and regional monitoring sites by season over the 18-yr period.....	3-1
Table 3-2. Summary of 24-hr PM_{10} concentrations ($\mu\text{g}/\text{m}^3$) at the Landfill, Community, and Landfill North monitoring sites and the South Coast AQMD Burbank, Santa Clarita, and Los Angeles regional sites when a federal PM_{10} exceedance ($>150\ \mu\text{g}/\text{m}^3$) occurred at either the Landfill or Community site.....	3-5
Table 3-3. Summary of state 24-hr PM_{10} exceedances (more than $50\ \mu\text{g}/\text{m}^3$) at the Landfill, Community, and Landfill North monitoring sites, and at the Burbank, Santa Clarita, and Los Angeles regional sites operated by the South Coast AQMD.....	3-10
Table 7-1. Sunshine Canyon Landfill monitoring site visits and field maintenance and operations in Year 18.	7-1
Table 7-2. Community monitoring site visits and field maintenance and operations in Year 18.	7-2

Acronyms

Term	Definition
BAM	Beta Attenuation Monitor
BC	black carbon
CAAQS	California ambient air quality standards
CARB	California Air Resources Board
Community site	Van Gogh Elementary School
DPM	diesel particulate matter
EPA	U.S. Environmental Protection Agency
FRM	Federal Reference Method
FSEIR	Final Supplemental Environmental Impact Report
HMS	Hazard Mapping System
ISCST3	Industrial Source Complex Short Term
Landfill site	Sunshine Canyon Landfill
LFG	landfill gas
MATES V	Multiple Air Toxics Exposure Study V
NAAQS	National Ambient Air Quality Standards
NWD	non-working days
NWH	non-working hours
PM ₁₀	particulate matter less than 10 microns in aerodynamic diameter
SoCAB	Southern California Air Basin
South Coast AQMD	South Coast Air Quality Management District
Spheros	Spheros Environmental
WD	working days
WH	working hours
WS/WD	wind speed/wind direction

Executive Summary

Continuous monitoring of particulate matter less than 10 microns in aerodynamic diameter (PM₁₀), black carbon (BC), wind speed, and wind direction began at the Sunshine Canyon Landfill (Landfill site) and at Van Gogh Elementary School (Community site) in Granada Hills, CA, in fall 2007. This Eighteenth Annual Report includes data summaries, analysis, and interpretation drawn from 18 complete years of data from the Landfill and Community monitoring sites, one year of data from a temporary Landfill North site that was installed upwind of the landfill for one year (2016), and data from the baseline year (November 22, 2001–November 21, 2002). These data are used to characterize ambient PM₁₀ and BC concentrations on a neighborhood scale in the context of the Southern California Air Basin (SoCAB), and to evaluate the impact of landfill operations on air quality in the community.

The following conclusions are based on data from 18 years of continuous monitoring of PM₁₀, BC, and meteorology at the Landfill and Community monitoring sites. Additionally, this report highlights 2025 (Year 18).

- Over the 18-yr monitoring period, the federal 24-hr PM₁₀ standard was exceeded on 59 occasions at the Landfill site, with a record number of 11 exceedances occurring in 2017 (Year 10). The federal standard was exceeded on a total of 6 occasions at the Community site in the 18-yr monitoring period. In 2025 (Year 18), the federal standard was exceeded on one occasion at the Landfill site and no occasions at the Community site.
- The more stringent state 24-hr PM₁₀ standard was exceeded on 18 occasions at the Landfill site in 2025 (Year 18), which is the lowest of any year on record. The Community site experienced four exceedances of the state standard in 2025 (Year 18), which is the same number of exceedances as in Year 16 and Year 17, and is at the lower end of the amount of annual state exceedances for the site.
- The Landfill site's PM₁₀ federal and state exceedances were accompanied by high wind speeds and wind direction that fell within a narrow sector encompassing the active portion of the landfill. State exceedance days at the Landfill site are also accompanied by low-speed winds from the Los Angeles basin (south and southeast), suggesting that the addition of elevated concentrations within the basin can push the Landfill site's PM₁₀ concentrations over the state threshold. On days when PM₁₀ concentrations exceeded the state standard at the Community site, wind speeds were relatively low and wind direction was predominantly from the Los Angeles basin (from the southeast). This indicates that regional contributions are the main driver of state PM₁₀ standard exceedances at the Community site.
- Monthly average PM₁₀ concentrations at the Landfill site have been higher than those recorded at the regional monitoring site in downtown Los Angeles since 2017 (Year 10). In Year 18, monthly average PM₁₀ concentrations were much lower than the previous five years, and the monthly average PM₁₀ concentrations did not exceed the state standard of 50 µg/m³.

Further, the annual average PM₁₀ concentration is below the state standard and exhibited a decrease from the previous years. The annual average PM₁₀ concentration at the Community site has been steadily decreasing since Year 7 and is now slightly lower than the annual average concentration measured at Santa Clarita in Year 18.

- To estimate landfill contributions to PM₁₀ and BC concentrations at both the Landfill and Community sites, we compared the difference between PM₁₀ and BC concentrations at each site under two wind sectors (“from landfill” and “from SoCAB”) and two working categories (non-working day/hour and working day/hour).
 - The greatest difference in PM₁₀ and BC concentrations between the Landfill and Community sites was observed during periods of highest activity levels (i.e., working hours on working days).
 - The Community site measured slightly higher PM₁₀ and BC concentrations on working days (both working hours and non-working hours) compared to non-working days (both working hours and non-working hours) when the wind was from the landfill. A similar workday/non-workday pattern exists for PM₁₀ concentrations at Burbank and Los Angeles regional sites, which may indicate that the increase is attributable to higher levels of emissions in general on working days relative to non-working days. However, PM₁₀ concentrations measured at the Community site were significantly lower than at regional monitoring sites and the Landfill site.
 - When the wind was coming from SoCAB, PM₁₀ values at the Landfill site were slightly higher than the Community site during non-working and working hours. On days in the highest activity-level category, regional contributions of PM₁₀ combined with local landfill contributions to increase PM₁₀ concentrations at the Landfill site. In 2025 (Year 18), PM₁₀ levels were slightly lower at the Community site than the Landfill site when wind was from SoCAB for all categories except working hours on non-working days.
 - When wind was from SoCAB, BC concentrations were higher at the Community site than the Landfill site during the working day/hour and non-working day/working hour categories, but slightly lower during the working day/non-working hour and non-working day/hour categories. This suggests that increased regional BC concentrations contributed to BC levels at the Community site.

1. Introduction

Two air quality monitoring sites were established by operators of the Sunshine Canyon Landfill in 2001. One monitoring site is on a high-elevation ridge on the southern edge of the Sunshine Canyon Landfill (Landfill site). The second site is at Van Gogh Elementary School in the nearby community of Granada Hills (Community site). These sites were established to monitor particulate matter less than 10 microns in aerodynamic diameter (PM₁₀), black carbon (BC) as a surrogate for diesel particulate matter (DPM), wind direction, and wind speed, in fulfillment of the stipulations set forth in the City of Los Angeles' Conditions of Approval for the expansion of the Sunshine Canyon Landfill in the City of Los Angeles (Section C.10.a of Ordinance No. 172,933). In 2009, the County of Los Angeles Department of Regional Planning and Public Works adopted conditions (County Condition 81) very similar to the City's conditions, governing ambient air quality monitoring for the County portion of the landfill.

1.1 Baseline Year and Continuous Monitoring

Continuous monitoring of PM₁₀, BC, and meteorology was performed during a baseline year between November 22, 2001, and November 21, 2002, and a report of the baseline year results was produced by ENVIRON International Corporation.¹ Between the time that the baseline studies were completed and November 2007, when continuous monitoring began, ambient sampling for PM₁₀, BC, and landfill gases (LFG) was planned at a nominal frequency of four times each year by ENVIRON International Corporation. Data from those years are not included in this report.

Beginning in 2007, ambient monitoring of particulate matter (and LFGs in some years) at the Landfill and Community sites became the responsibility of Spheros Environmental (Spheros, formerly Sonoma Technology). Spheros's technical approach to monitoring PM₁₀ and BC was based on continuous monitoring (hourly, year-round), whereas previous monitoring was limited to four events per year. Continuous year-round monitoring of PM₁₀ and BC allows greater potential to evaluate times when air flows from the landfill to the Community site, as well as to evaluate diurnal trends, day-of-week differences, seasonal differences, and annual trends in pollutant concentrations compared to regional monitors operated by the South Coast Air Quality Management District (South Coast AQMD) and the California Air Resources Board (CARB).

November 21, 2025, marked the completion of 18 full years of continuous monitoring of PM₁₀, BC, and meteorology at the two main monitoring locations. Data capture rates and the quality of the captured data have generally been very high. A few discrete events have interrupted data capture at one or both sites. For example, the Sayre Fire in late 2008 took out power at the Landfill site for several

¹ ENVIRON International Corporation (2003) Results of the baseline ambient air monitoring program for the Sunshine Canyon Landfill. Final report prepared for Browning-Ferris Industries of California, Inc., by ENVIRON International Corporation, Contract No. 03-9660A, June 6.

weeks. Monitoring equipment upgrades in 2010 caused some loss of data because instruments were temporarily removed. There was significant loss of PM₁₀ data during the fourth quarter of Years 9 and 11 because the Beta Attenuation Monitor (BAM) instruments were removed from the field and sent to the manufacturer for maintenance. In 2019, the Landfill site was without power for about half a month in October due to the Saddle Ridge Fire. During the Year 13 spring quarter, about a third of the captured PM₁₀ data were invalidated due to limited access to sites for scheduled instrument maintenance during the COVID-19 shelter-in-place order. Even with these interruptions, however, PM₁₀ data completeness statistics for the 18 years indicate average data capture rates of approximately 97% at the Landfill site and the Community site (see [Section 2](#)). On average, approximately 4% of all captured data at the Landfill and Community sites were judged as invalid.

1.2 Report Overview

In this report, the high-quality, high-time-resolution data captured over the 18 years between November 2007 and November 2025 at the Landfill and Community sites are analyzed and summarized to offer a realistic characterization of ambient air quality concentrations at the Sunshine Canyon Landfill and the Granada Hills community, and to provide perspective on air quality at the landfill and the local community in the context of the greater South Coast Air Basin (SoCAB).

- [Section 2](#) of this report discusses data completeness.
- [Section 3](#) covers PM₁₀ exceedances of state and federal standards.
- [Section 4](#) discusses regional comparisons of PM₁₀ concentrations.
- [Section 5](#) describes the effects of wind direction and work activity levels on PM₁₀ and BC concentrations at the Landfill and Community sites.
- [Section 6](#) discusses the landfill's impact on ambient PM₁₀ and BC concentrations.
- [Section 7](#) describes routine field operations and recent upgrades to site infrastructure.
- [Appendix A](#) discusses regional comparisons of BC with data from the most recent Multiple Air Toxics Exposure Study V (MATES V).
- Additional analyses of wind and the Landfill North site data are provided in [Appendix B](#).
- [Appendix C](#) compares the Environmental Impact Report's estimated annual increment from landfill emissions to ambient air toxics concentrations and ambient air toxics from the 2016-2017 measurements made at the Landfill and Community sites.

Regulatory standards for pollutants are commonly used to judge the compliance status of air quality management districts. Currently, the only federal health-based standard for PM₁₀ is the daily (24-hr) average concentration of 150 µg/m³. The State of California's PM₁₀ 24-hr standard (50 µg/m³) is more stringent than the federal standard. In this report, both the 24-hr federal standard and the 24-

hr state standard are used as a benchmark metric for evaluating the specific monitoring locations in relation to each other and to the standards.

Regional comparisons of ambient PM₁₀ concentrations are used to place the Landfill and Community sites within the larger context of regional concentrations. For these comparisons, three of the closest regional monitoring sites, operated by South Coast AQMD, were chosen: downtown Los Angeles (North Main Street), Burbank (West Palm),² and Santa Clarita. **Figure 1-1** shows the relative locations of the sites.




Figure 1-1. Locations of the Landfill and Community sites in relation to the three South Coast AQMD PM₁₀ sites and four MATES V BC sites used for regional comparisons. The Landfill site is labeled “Landfill” and the Community site is labeled “Community.” In MATES V documentation, the Los Angeles North Main site is referred to as “Central LA.” The Landfill North site was operated from 2016–2017 and is not shown here.

² PM₁₀ monitoring at the Burbank (West Palm) site was discontinued in July 2014.

Ambient concentrations of BC as a surrogate for DPM continue to receive increased interest statewide, nationally, and globally. South Coast AQMD has shown that DPM is one of the primary air toxics of concern in the SoCAB. To place the Landfill and Community sites within the larger context of regional concentrations, four of the closest regional monitoring sites from the Multiple Air Toxics Exposure Study (MATES V, May 2018–April 2019),³ also operated by the South Coast AQMD, were selected: Burbank (approximately the same location as the Burbank PM₁₀ site), Central LA (i.e., Los Angeles North Main) (approximately the same location as the Los Angeles PM₁₀ site), Huntington Park, and Pico Rivera. Note that this regional comparison spans only the 1-yr study period of the MATES V study (Appendix A).

1.3 Methods and Operations Background

Aethalometers measure BC concentrations using an optical attenuation technique, and measurements are subject to what is known as a tape saturation effect, where the buildup of BC on the tape causes an artifact affecting the accuracy of the measured concentration.^{4,5} Instrument response is dampened with heavier loading (i.e., higher concentrations) of BC particles on the tape. This artifact can bias reported BC concentrations low. However, mathematical methods to correct the BC concentrations are available and are widely used. BC values from the Landfill and Community sites were compensated for this tape saturation effect and therefore are representations of ambient concentrations.

Meteorological factors and landfill work activity levels are known to have an impact on local and regional pollutant concentrations. An analysis based on wind direction and landfill working versus non-working days and hours is used to quantify the relationship of these factors to PM₁₀ and BC concentrations. This analysis also provides quantitative estimates of landfill contributions to ambient concentrations of PM₁₀ and BC. A summary of the analytical method is presented in Section 6, with additional analyses in Appendix B.

³ Information at <http://www.aqmd.gov/home/air-quality/air-quality-studies/health-studies/mates-v>.

⁴ Drinovec L et al. (2014) The "dual-spot" Aethalometer: an improved measurement of aerosol black carbon with real-time loading compensation. *Atmos. Meas. Tech. Discuss.*, 7(9), 10179–10220, doi: 10.5194/amtd-7-10179-2014. Available at <http://www.atmos-meas-tech-discuss.net/7/10179/2014/>.

⁵ Allen G. (2014) Analysis of spatial and temporal trends of black carbon in Boston. Report prepared by Northeast States for Coordinated Air Use Management (NESCAUM), Boston, MA, January. Available at nescaum.org/documents/analysis-of-spatial-and-temporal-trends-of-black-carbon-in-boston/nescaum-boston-bc-final-rept-2014.pdf.

2. Data Completeness

Table 2-1 shows completeness statistics for all measured variables for the 18 years considered in this analysis. Percent Data Capture is the percent of hourly data values that were collected divided by the total number of expected data intervals in the date range (e.g., 24 hourly data values are expected per day, and 8,760 hourly data values are expected per year—8,784 during leap years). Percent Data Valid or Suspect is the percent of data values that are either valid or suspect divided by the number of captured data values. Percent Data Suspect is the percentage of data values that are labeled as suspect divided by the number of captured data values. WS/WD is wind speed/wind direction.

Except for Year 2 (when the Sayre Fire shut down the Landfill site's data collection effort from November 15, 2008, through January 8, 2009) and Years 9 and 11 (due to instrument maintenance), the percent data capture for PM₁₀ exceeded 90% in each site-year at both the Landfill and Community sites. The percent data capture averaged 96.4% at the Landfill site and 98.4% at the Community site over all 18 years. The percent data capture for PM₁₀ in the 18th year is 99.6% at the Landfill site and 97.3% at the Community site. The percent data capture for BC at the Landfill site averaged 94.5% over 18 years and was 99.9% in the 18th year and averaged 95.8% over 18 years and was 99.9% in the 18th year at the Community site.

As shown in Table 2-1, the percent data capture for WS/WD was 100% at the Landfill site in Year 18 and averaged 96.9% over all 18 years. At the Community site in Year 10, data logging computer failures caused significant WS/WD data loss, resulting in roughly 65% data capture. In Year 18 at the Community site, the percent data capture for WS/WD was 98.9%, and over the 18-yr period the data capture at the Community site averaged 96.8%.

Table 2-1. Data completeness statistics for hourly data during Years 1-18 of continuous monitoring and overall 18-yr averages. The beginning and end dates for each year are chosen to allow comparison with the baseline year (Nov. 22, 2001-Nov. 21, 2002).

Years	Monitoring Location	Percent Data Capture (%)			Percent Data Valid or Suspect (%)			Percent Data Suspect (%)		
		PM ₁₀	BC	WS/WD	PM ₁₀	BC	WS/WD	PM ₁₀	BC	WS/WD
Yr. 1 Nov. 22, 2007– Nov. 21, 2008	Sunshine Canyon Landfill Site	94.2%	90.7%	88.3%	98.0%	99.9%	93.3%	0.0%	0.0%	0.0%
	Van Gogh Elementary School Site	95.8%	92.3%	95.4%	96.0%	100.0%	94.7%	0.1%	0.0%	0.0%
Yr. 2 Nov. 22, 2008– Nov. 21, 2009	Sunshine Canyon Landfill Site	86.6%	81.3%	86.8%	97.9%	100.0%	98.3%	0.0%	0.0%	0.0%
	Van Gogh Elementary School Site	98.7%	98.5%	99.9%	96.3%	100.0%	99.9%	0.0%	0.0%	0.0%
Yr. 3 Nov. 22, 2009– Nov. 21, 2010	Sunshine Canyon Landfill Site	99.7%	87.8%	98.4%	98.2%	100.0%	99.2%	0.1%	0.0%	4.3%
	Van Gogh Elementary School Site	98.4%	87.9%	98.3%	97.0%	100.0%	100.0%	0.5%	23.3% ^a	0.0%
Yr. 4 Nov. 22, 2010– Nov. 21, 2011	Sunshine Canyon Landfill Site	90.8%	99.6%	99.9%	96.9%	100.0%	97.5%	0.3%	0.0%	1.6%
	Van Gogh Elementary School Site	100.0%	99.8%	100.0%	99.2%	99.9%	96.3%	0.5%	0.0%	0.0%
Yr. 5 Nov. 22, 2011– Nov. 21, 2012	Sunshine Canyon Landfill Site	99.1%	99.6%	99.4%	95.4%	99.9%	96.7%	5.4%	0.0%	1.0%
	Van Gogh Elementary School Site	94.1%	99.9%	98.7%	98.1%	99.9%	96.1%	0.2%	0.0%	0.0%
Yr. 6 Nov. 22, 2012– Nov. 21, 2013	Sunshine Canyon Landfill Site	99.9%	99.7%	98.7%	98.6%	99.9%	100.0%	0.8%	0.0%	0.0%
	Van Gogh Elementary School Site	100.0%	99.8%	99.4%	97.7%	100.0%	100.0%	0.4%	0.1%	0.0%
Yr. 7 Nov. 22, 2013– Nov. 21, 2014	Sunshine Canyon Landfill Site	100.0%	87.9%	98.1%	99.3%	100.0%	100.0%	0.3%	0.0%	0.0%
	Van Gogh Elementary School Site	100.0%	99.1%	98.5%	98.0%	100.0%	100.0%	0.2%	0.6%	0.0%

Years	Monitoring Location	Percent Data Capture (%)			Percent Data Valid or Suspect (%)			Percent Data Suspect (%)		
		PM ₁₀	BC	WS/WD	PM ₁₀	BC	WS/WD	PM ₁₀	BC	WS/WD
Yr. 8 Nov. 22, 2014– Nov. 21, 2015	Sunshine Canyon Landfill Site	99.9%	88.4%	98.6%	98.3%	100.0%	100.0%	0.3%	0.1%	0.0%
	Van Gogh Elementary School Site	99.9%	85.1%	99.0%	82.2%	100.0%	100.0%	0.1%	0.0%	0.0%
Yr. 9 Nov. 22, 2015– Nov. 21, 2016	Sunshine Canyon Landfill Site	91.8%	93.3%	99.16%	81.3%	99.8%	100.0%	0.0%	8.7%	0.0%
	Van Gogh Elementary School Site	89.9%	92.4%	99.18%	89.1%	99.7%	100.0%	0.0%	0.3%	0.0%
	Sunshine Canyon Landfill North Site ^b	80.3%	85.6%	88.0%	94.8%	99.9%	100.0%	0.0%	0.2%	0.0%
Yr. 10 Nov. 22, 2016– Nov. 21, 2017	Sunshine Canyon Landfill Site	98.6%	94.0%	97.5%	99.1%	100.0%	100.0%	0.0%	0.0%	0.0%
	Van Gogh Elementary School Site	99.9%	91.5%	64.7%	99.8%	99.8%	100.0%	0.0%	0.0%	0.0%
	Sunshine Canyon Landfill North Site ^b	99.6%	90.3%	99.6%	99.4%	100.0%	100.0%	0.0%	17.5%	0.0%
Yr. 11 Nov. 22, 2017– Nov. 21, 2018	Sunshine Canyon Landfill Site	87.0%	91.1%	91.6%	98.1%	100.0%	100.0%	0.0%	10.9%	0.0%
	Van Gogh Elementary School Site	100.0%	95.6%	100.0%	98.0%	99.3%	100.0%	0.0%	6.9%	0.0%
Yr. 12 Nov. 22, 2018– Nov. 21, 2019	Sunshine Canyon Landfill Site	95.9%	95.0%	95.8%	98.4%	100.0%	94.6%	0.1%	2.2%	0.0%
	Van Gogh Elementary School Site	99.9%	96.3%	100.0%	97.1%	99.8%	99.9%	0.0%	4.1%	0.0%
Yr. 13 Nov. 22, 2019– Nov. 21, 2020	Sunshine Canyon Landfill Site	99.7%	99.6%	99.6%	94.4%	100.0%	99.2%	0.0%	3.6%	0.0%
	Van Gogh Elementary School Site	99.7%	99.1%	99.1%	90.4%	100.0%	99.2%	0.0%	3.7%	0.0%
Yr. 14 Nov. 22, 2020– Nov. 21, 2021	Sunshine Canyon Landfill Site	97.8%	97.6%	95.7%	93.9%	100.0%	100.0%	0.0%	3.2%	0.0%
	Van Gogh Elementary School Site	98.2%	95.1%	99.0%	97.2%	100.0%	100.0%	0.0%	8.0%	0.0%

Years	Monitoring Location	Percent Data Capture (%)			Percent Data Valid or Suspect (%)			Percent Data Suspect (%)		
		PM ₁₀	BC	WS/WD	PM ₁₀	BC	WS/WD	PM ₁₀	BC	WS/WD
Yr. 15 Nov. 22, 2021– Nov. 21, 2022	Sunshine Canyon Landfill Site	98.4%	99.8%	98.7%	94.5%	100.0%	100.0%	0.0%	3.8%	0.0%
	Van Gogh Elementary School Site	99.7%	99.3%	98.9%	98.3%	100.0%	100.0%	5.7%	4.9%	0.0%
Yr. 16 Nov. 22, 2022– Nov. 21, 2023	Sunshine Canyon Landfill Site	95.7%	99.1%	98.3%	95.6%	98.4%	100.0%	58.0% ^c	2.9%	0.0%
	Van Gogh Elementary School Site	99.3%	98.9%	97.2%	99.8%	100.0%	100.0%	9.2%	4.0%	0.0%
Yr. 17 Nov. 22, 2023– Nov. 21, 2024	Sunshine Canyon Landfill Site	96.7%	96.2%	99.6%	98.2%	99.9%	100.0%	27.8%	0.3%	11.1%
	Van Gogh Elementary School Site	99.7%	99.0%	100.0%	99.8%	99.9%	100.0%	13.8%	0.3%	0.0%
Yr. 18 Nov. 22, 2024– Nov. 21, 2025	Sunshine Canyon Landfill Site	99.6%	99.9%	100.0%	97.0%	99.8%	100.0%	25.4%	0.1%	0.2%
	Van Gogh Elementary School Site	97.3%	99.9%	98.9%	97.5%	99.3%	100.0%	11.2%	1.7%	0.0%
Eighteen-Yr. Average	Sunshine Canyon Landfill Site	96.4%	94.5%	96.9%	96.0%	99.9%	98.9%	6.6%	2.0%	0.8%
	Van Gogh Elementary School Site	98.4%	95.8%	96.8%	96.1%	99.9%	99.2%	1.8%	3.3%	0.0%

^a Three-fourths of the data from the June 2010–August 2010 quarter were suspect because flow rates as measured by the reference flow meter were outside of tolerance levels. This was due to a leak in the push-to-connect fitting at the back of the Aethalometer. Further details can be found in the Eleventh Quarterly report. This quarter negatively affects the 18-yr average for percent suspect. Without this quarter, the 18-yr average at the Community monitoring site would reduce from 3.3% to 2.1%.

^b Sunshine Canyon Landfill North site was operated from June 2016 through May 2017.

^c The BAM analyzer at the Landfill site experienced multiple issues throughout the 2023 winter, spring, and fall quarters that compromised data collection operations and data quality. An air conditioning unit on the trailer that houses the BAM experienced multiple malfunctions, winter storms caused power outages, and water from heavy rains and odor misting saturated the BAM sampling inlet, condensation jar, and sampling lines. These conditions led to a high abundance of suspect data. Without this quarter, the 18-yr average would be 3.6% instead of 6.6%.

3. PM₁₀ Exceedances

The Clean Air Act requires the U.S. Environmental Protection Agency (EPA) to set National Ambient Air Quality Standards (NAAQS, 40 CFR Part 50) for pollutants considered harmful to public health and the environment.⁶ PM₁₀ is included in the NAAQS. Currently, the only federal health-based standard for PM₁₀ is the daily (24-hr) average concentration of 150 µg/m³.

In 1959, California enacted legislation requiring the state Department of Public Health to establish air quality standards and necessary controls for motor vehicle emissions.⁷ California law continues to mandate California ambient air quality standards (CAAQS), which are often more stringent than national standards. The State of California's current 24-hr standard for PM₁₀ is 50 µg/m³. **Table 3-1** shows the number of federal and state PM₁₀ exceedances for the Landfill, Community, and regional monitoring sites by season over the 18-yr period. Additional information on the federal and state exceedance days are described in the following sections.

Table 3-1. Total number of 24-hr federal (150 µg/m³) and state (50 µg/m³) PM₁₀ exceedances at the Landfill, Community, and regional monitoring sites by season over the 18-yr period.

Exceedance Type	Season ^a	Sunshine Canyon Landfill	Community Site	Burbank ^b	Los Angeles North Main	Santa Clarita ^c
# of Federal Exceedances	Spring	16	1	0	0	0
	Summer	2	0	0	1	0
	Fall	25	5	0	1	0
	Winter	16	0	0	1	0
Total # of Federal Exceedances		59	6	0	3	0
# of State Exceedances	Spring	318	68	6	116	2
	Summer	481	128	6	114	1
	Fall	415	94	9	220	6
	Winter	186	34	7	173	2
Total # of State Exceedances		1,400	324	28	623	11

Notes:

^a Spring: March 1–May 31; Summer: June 1–August 31; Fall: Sept. 1–Nov. 30; Winter: Dec. 1–Feb. 28 (Feb. 29 for leap year).

^b The Burbank site was discontinued in July 2014.

^c Samples from the Santa Clarita site are collected every six days.

⁶ <https://www.epa.gov/criteria-air-pollutants/naaqs-table>.

⁷ <https://www.arb.ca.gov/research/aaqs/caaqs/caaqs.htm>.

3.1 Federal Exceedances

Figure 3-1 depicts the number of federal 24-hr PM_{10} exceedances measured at the Landfill, Community, and regional monitoring sites for each year of the 18-yr period. In Year 18, the federal standard was exceeded on one occasion at the Landfill site and was not exceeded at the Community site.




Figure 3-1. The number of federal exceedances of 24-hr PM_{10} at the Landfill, Community, and regional monitoring sites by year over the 18-yr study period. (e.g., Monitoring Year 1 refers to Nov. 22, 2007–Nov. 21, 2008).

The federal 24-hr PM_{10} standard (i.e., $150 \mu\text{g}/\text{m}^3$) was exceeded at the Landfill site on January 7, 2025, but not at the Community site. **Figure 3-2** shows 24-hr PM_{10} concentrations at sites across the greater Los Angeles area on January 7, 2025, with Hazard Mapping System (HMS) fire and smoke products and January 2025 fire perimeters overlaid. **Figure 3-3** shows a time series of 1-hour PM_{10} , BC, and wind data recorded five days before and after the January 7, 2025, PM_{10} exceedance.

PM_{10} concentrations in the Los Angeles area were elevated on January 7, 2025. At the Landfill site, high PM_{10} concentrations on January 7 are likely due mostly to wind-blown dust, but may have also been elevated by nearby fire activity. As shown by the HMS fire detections (red triangles) in the map, there were three major active fires on January 7, including the Palisades Fire, Eaton Fire, and Hurst Fire, near the Sunshine Canyon Landfill. Each of the three fires began on January 7 and spread rapidly due to high winds and dry conditions. The Palisades Fire, Eaton Fire, and Hurst Fire were detected at approximately 10:00 a.m. PST, 6:30 p.m. PST, and 10:00 p.m. PST, respectively. Of the three actively burning fires on January 7, the Hurst Fire may have contributed to elevated PM_{10} and BC.

concentrations at both the Landfill site and Community site late in the evening on January 7 and into January 8 as the fire spread rapidly. As shown in Figure 3-3, PM₁₀ concentrations at both sites began increasing early in the morning on January 7, reached a peak late in the evening, then began to decrease to baseline concentrations into January 8 and 9. During this increase in concentrations, wind direction at both sites shifted abruptly to between approximately 0 and 45°, which is from the direction of the Hurst Fire relative to the sites. In addition, BC concentrations at both sites sharply increased at the end of January 7 and into January 8 as smoke moved into the area. However, the elevated PM₁₀ concentrations throughout the day on January 7, which led to the exceedance, were likely caused in part by wind-blown dust rather than smoke impacts because the Hurst Fire began late in the evening on January 7. Further, wind speeds at the Landfill site were elevated in the days before and the day of January 7, 2025, with the PM₁₀ exceedance likely caused wind-blown dust that contributed to elevated PM₁₀ concentrations. On January 7, increasing wind speeds coincided with increasing PM₁₀ concentrations at the Landfill site. Hourly PM₁₀ concentrations and wind speed at the Landfill site reached a maximum of 549 µg/m³ and 44 mph, respectively, at 10:00 p.m. PST on January 7.




Figure 3-2. PM₁₀ concentrations at air monitoring sites across the Los Angeles area on January 7, 2025. Colors correspond to 24-hr PM₁₀ concentrations in µg/m³. HMS fire and smoke layers from January 7, 2025, the day of the exceedance, and fire perimeters from January 7, 2025, are overlaid. *Note: only the Palisades, Hurst, and Eaton fires were actively burning on January 7, 2025.*




Figure 3-3. Hourly time series of PM_{10} and BC concentrations (top two rows) and WS/WD measurements (bottom two rows) recorded at the Landfill and Community sites five days before and after the January 7, 2025, PM_{10} exceedance at the Landfill site. The gray rectangle indicates January 7, 2025.

Table 3-2 lists all the days during the past 18 years of continuous monitoring on which the federal 24-hr PM_{10} standard was exceeded at either the Landfill site or the Community site, along with 24-hr average concentrations from those days at the three comparative South Coast AQMD sites (Burbank, Santa Clarita, and downtown Los Angeles). The Burbank and Los Angeles sites have continuous (hourly) PM_{10} monitors, like those at the Landfill and Community sites. The Santa Clarita site, however, employs Federal Reference Method (FRM) sampling (integrated 24-hr samples collected on filters) on a one-in-six-day schedule. Note that a 75% data completeness threshold was used when calculating

the 24-hr average PM₁₀ from the hourly measurements. For example, 18 out of 24 data points (per day) need to be valid to calculate the 24-hr average value at a site with hourly measurements.

Table 3-2. Summary of 24-hr PM₁₀ concentrations (µg/m³) at the Landfill, Community, and Landfill North monitoring sites and the South Coast AQMD Burbank, Santa Clarita, and Los Angeles regional sites when a federal PM₁₀ exceedance (>150 µg/m³) occurred at either the Landfill or Community site.

Date	Landfill Site	Community Site	Landfill North Site	Burbank West Palm	Los Angeles Main Street	Santa Clarita
2/14/2008	167	48	-- b	19	30	-- b
5/21/2008	290	152	-- b	119	140	-- b
10/9/2008	158	104	-- b	--b	59	91
1/9/2009	185	71	-- b	-- b	68	-- b
5/6/2009	257	91	-- b	-- b	49	-- b
10/27/2009	239	165	-- b	130	147	-- b
1/20/2011	207	28	-- b	26	46	-- b
4/30/2011	221	32	-- b	25	40	-- b
11/2/2011	263	43	-- b	37	56	-- b
5/22/2012	186	61	-- b	34	-- a	-- b
10/26/2012	227	49	-- b	31	40	-- b
3/21/2013	181	34	-- b	32	37	-- b
4/8/2013	174	64	-- b	53	-- b	-- b
10/4/2013	200	64	-- b	28	58	-- b
12/4/2013	155	18	-- b	21	-- a	-- b
12/9/2013	181	31	-- b	24	34	-- b
7/22/2016	183	51	66	-- c	53	-- b
7/30/2016	153	129	209	-- c	36	-- b
11/17/2016	178	38	-- b	-- c	51	-- b
12/2/2016	245	76	84	-- c	35	22
12/18/2016	204	32	21	-- c	26	-- b
3/27/2017	170	37	26	-- c	28	-- b
4/20/2017	236	37	30	-- c	35	-- b
4/21/2017	167	41	29	-- c	40	-- b
4/25/2017	191	42	38	-- c	28	67

Date	Landfill Site	Community Site	Landfill North Site	Burbank West Palm	Los Angeles Main Street	Santa Clarita
4/27/2017	184	45	45	-- c	45	-- b
4/28/2017	165	47	46	-- c	33	-- b
10/9/2017	200	61	-- b	-- c	61	-- b
10/24/2017	276	35	-- b	-- c	39	-- b
11/21/2017	170	30	-- b	-- c	48	25
12/5/2017	225	62	-- b	-- c	54	-- b
12/17/2017	210	54	-- b	-- c	36	-- b
4/12/2018	237	50	-- b	-- c	40	-- b
11/8/2018	231	60	-- b	-- c	49	-- b
4/9/2019	193	53	-- b	-- c	59	42
10/10/2019	-- a	161	-- b	-- c	47	-- b
10/11/2019	-- a	259	-- b	-- c	45	-- b
10/25/2019	202	64	-- b	-- c	47	-- b
10/30/2019	-- a	324	-- b	-- c	64	63
10/31/2019	170	61	-- b	-- c	42	-- b
11/16/2019	157	30	-- b	-- c	43	-- b
11/25/2019	212	90	-- b	-- c	33	-- b
11/26/2019	188	123	-- b	-- c	94	-- b
12/17/2019	157	33	-- b	-- c	29	8
9/9/2020	157	51	-- b	-- c	48	-- b
10/16/2020	204	48	-- b	-- c	51	-- b
10/26/2020	204	111	-- b	-- c	185	-- b
11/26/2020	172	33	-- b	-- c	43	-- b
12/3/2020	235	53	-- b	-- c	57	-- b
12/7/2020	155	41	-- b	-- c	66	-- b
12/23/2020	173	44	-- b	-- c	54	35
1/19/2021	227	70	-- b	-- c	64	-- b
9/23/2021	152	22	-- b	-- c	36	-- b
10/11/2021	288	153	-- b	-- c	139	-- b
11/25/2021	262	37	-- b	-- c	46	-- b
4/5/2022	224	35	-- b	-- c	44	-- b

Date	Landfill Site	Community Site	Landfill North Site	Burbank West Palm	Los Angeles Main Street	Santa Clarita
11/16/2022	284	32	-- b	-- c	43	-- b
4/19/2023	154	34	-- b	-- c	-- b	-- b
10/29/2023	198	62	-- b	-- c	43	-- b
10/30/2023	239	46	-- b	-- c	40	-- b
12/9/2023	254	53	-- b	-- c	34	-- b
1/7/2025	163	64	-- b	-- c	123	81

Notes:

^a Not enough hourly data to meet the 75% data threshold standard.

^b No data available.

^c PM₁₀ monitoring at the Burbank West Palm regional site was discontinued in July 2014.

Over the past 18 years of continuous monitoring, the federal standard was exceeded on 59 occasions at the Landfill site. Of the 59 exceedances at the Landfill site, the Community site registered an exceedance on 3 of those days. While the South Coast AQMD sites in Burbank, Santa Clarita, and Los Angeles did not report exceedances on these three days, the 24-hr PM₁₀ concentrations were relatively high. The elevated concentrations at these sites suggest that, when regional concentrations are high, the combination of landfill and regional contributions can push the Community site's PM₁₀ concentrations over the federal standard. Additionally, the Community site had three federal exceedances in October 2019, likely due to wildfire activities (see the Twelfth Annual Report). However, over 18 years of monitoring, high regional concentrations combined with high landfill concentrations have only occurred on four days (05/21/2008, 10/27/2009, 10/26/2020, and 10/11/2021). More conclusive, however, is the insignificant effect on Community PM₁₀ concentrations that occur when Landfill concentrations exceed federal limits and regional concentrations are relatively low. As shown in Table 3-2 and Figure 3-1, this is particularly evident on the 11 federal exceedance days at the Landfill site in Year 10 (2017).

The Landfill site PM₁₀ federal exceedances listed in Table 3-2 were generally accompanied by high wind speeds, with wind direction falling within a narrow sector that encompasses the active portion of the landfill. Wind data from the Landfill site for all federal exceedance days are plotted in [Figure 3-4](#). A wind rose depicts how wind speed and direction are typically distributed at a particular location. Presented in a circular format, the length of each "spoke" is related to the frequency of time that wind blows from that direction. The color of each spoke indicates differences in wind speed. Most of the winds were from the northwest, passing directly over working areas of the landfill. Wind speeds were highest when the wind direction was from the northwest and north. Also shown in Figure 3-4 is wind data from the Community site for the six federal exceedance days. While the wind direction is also mainly from the north-northwest, wind speeds are significantly lower.




Figure 3-4. Wind rose from federal exceedance days during 18 continuous monitoring years at the Landfill (top right) and Community (bottom right) sites. Wind data at the Community site are replaced with those from the Reseda site from Year 11 to Year 16.

After 18 years of continuous data collection, it is clear that PM_{10} federal exceedances at the Landfill site are more common than they are in the Community or at regional monitoring sites, suggesting that surface material is being entrained at high wind speeds and subsequently detected by the Landfill monitoring site. By the time these air parcels reach the Community or regional sites, they have been diluted, and some of the larger particles may have been removed by deposition. Additionally, because of the relatively large elevation change and short distance between the Landfill and Community monitoring sites, air parcels under high-speed winds impacting the Landfill monitoring site can travel over the Community monitoring site.

As mentioned above, without the existence of an upwind monitoring site (located north-northwest of the existing Landfill monitoring site, on the north end of the landfill activity area), quantifiable air pollutant contributions from landfill activities cannot be determined.

3.2 State Exceedances

Figure 3-5 depicts the number of 24-hr PM_{10} California state exceedances measured at the Landfill, Community, and regional monitoring sites for each year of the 18-yr period. State exceedances are more common across sites than the federal exceedances shown in Figure 3-1. Although state exceedances at the Landfill site declined three years in a row between 2017 (Year 10) and 2019 (Year 12), 2020 (Year 13) saw a large increase as PM_{10} concentrations exceeded the state standard 180 times. The following year (Year 14) recorded 216 exceedances of the state standard, a record number. 2022 (Year 15) had the second highest number of state exceedances, 189, over the 16-yr monitoring period. In 2023 (Year 16), the number of state exceedances declined to 127. In Year 17, the number of state exceedances declined further to 22. In Year 18, the number of state exceedances declined again to a record low of 18.




Figure 3-5. Number of state exceedances of 24-hr PM_{10} at the Landfill, Community, and regional monitoring sites by year over the 18-yr study period. (e.g., Monitoring Year 1 refers to Nov. 22, 2007–Nov. 21, 2008).

The Community site has seen a low number (< 10) of state 24-hr PM_{10} standard exceedances consistently since 2015. However, in Year 12, the number of exceedances at the Community site rose to 16 due to regional and state wildfire and smoke activities in the fall quarter of 2019. In Year 15, state exceedances at the Community site decreased to one, setting the record for the lowest number of state exceedances over the 18-yr monitoring period. In Year 16, state exceedances at the Community site increased slightly to four, which is the second-lowest number of state exceedances

at the Community site over the 18-yr monitoring period. The number of state exceedances (4) at the Community site in Year 17 and Year 18 remained constant from Year 16.

Table 3-3 lists the number of days during the past 18 years of continuous monitoring when the state 24-hr PM₁₀ standard was exceeded at all three monitoring sites operated by Spheros, along with the three comparative South Coast AQMD sites (Burbank, Santa Clarita, and downtown Los Angeles).

Table 3-3. Summary of state 24-hr PM₁₀ exceedances (more than 50 µg/m³) at the Landfill, Community, and Landfill North monitoring sites, and at the Burbank, Santa Clarita, and Los Angeles regional sites operated by the South Coast AQMD.

Site	Year	No. of Exceedances	No. of Valid 24-hr Averages	% Exceedances
Sunshine Canyon Landfill	Year 1	70	337	21%
	Year 2	57	307	19%
	Year 3	32	354	9%
	Year 4	57	320	18%
	Year 5	47	341	14%
	Year 6	64	359	18%
	Year 7	57	364	16%
	Year 8	34	358	9%
	Year 9	32	270	12%
	Year 10	98	356	28%
	Year 11	52	311	17%
	Year 12	48	342	14%
	Year 13	180	343	52%
	Year 14	216	331	65%
	Year 15	189	337	56%
	Year 16	127	332	38%
	Year 17	22	343	6%
	Year 18	18	352	5%
Sunshine Canyon Landfill North ¹	Year 9	77	275	28%
	Year 10	14	190	7%
Community site	Year 1	44	335	13%
	Year 2	48	341	14%

Site	Year	No. of Exceedances	No. of Valid 24-hr Averages	% Exceedances
	Year 3	41	346	12%
	Year 4	26	362	7%
	Year 5	25	336	7%
	Year 6	38	354	11%
	Year 7	29	359	8%
	Year 8	5	299	2%
	Year 9	5	291	2%
	Year 10	8	365	2%
	Year 11	7	357	2%
	Year 12	16	354	5%
	Year 13	14	329	4%
	Year 14	5	347	1%
	Year 15	1	356	< 1%
	Year 16	4	362	1%
	Year 17	4	365	1%
	Year 18	4	348	1%
Burbank ²	Year 1	12	217	6%
	Year 2	5	58	9%
	Year 3	5	363	1%
	Year 4	0	362	0%
	Year 5	4	366	1%
	Year 6	1	360	0%
	Year 7	1	200	1%
Los Angeles North Main	Year 1	108	312	35%
	Year 2	113	354	32%
	Year 3	25	342	7%
	Year 4	25	317	8%
	Year 5	39	335	12%
	Year 6	32	332	9%
	Year 7	43	342	13%
	Year 8	35	335	10%

Site	Year	No. of Exceedances	No. of Valid 24-hr Averages	% Exceedances
	Year 9	30	360	8%
	Year 10	28	338	8%
	Year 11	48	364	13%
	Year 12	16	358	4%
	Year 13	31	360	9%
	Year 14	17	357	5%
	Year 15	11	362	3%
	Year 16	3	357	<1%
	Year 17	6	360	2%
	Year 18	13	360	4%
Santa Clarita ³	Year 1	2	56	4%
	Year 2	1	53	2%
	Year 3	0	57	0%
	Year 4	0	56	0%
	Year 5	0	55	0%
	Year 6	0	60	0%
	Year 7	0	59	0%
	Year 8	0	53	0%
	Year 9	1	60	2%
	Year 10	1	57	2%
	Year 11	1	50	2%
	Year 12	1	60	2%
	Year 13	1	40	3%
	Year 14	0	50	0%
	Year 15	0	61	0%
	Year 16	0	55	0%
	Year 17	1	61	2%
	Year 18	1	52	2%

Notes:

1. Sunshine Canyon Landfill North Site operated June 2016 through May 31, 2017.
2. PM₁₀ monitoring was discontinued in July 2014 at the Burbank site.
3. FRM sampling (integrated 24-hr samples on filters) on a one-in-six-day schedule at the Santa Clarita site.

Similar to the federal exceedance pattern (discussed in Section 3.1), the Landfill PM_{10} state exceedances were accompanied by high wind speeds, with wind direction falling within a narrow sector that encompassed the active portion of the landfill. However, as shown in Figure 3-6, state exceedance days at the Landfill site were also accompanied by low wind speeds and wind directions from the Los Angeles basin (south and southeast). These elevated concentrations within the basin, in combination with landfill contributions, can push the Landfill site's PM_{10} concentrations over the state standard. To help explain this pattern and to emphasize the importance of meteorology's effect on measured pollutant levels, the Ninth Annual Report provided meteorological data measured at the Landfill site for the years 2008 through 2016. These data demonstrated that measurements at the Landfill site are dominated by summer season wind flow from the south to south-southeast, and thus by regional PM_{10} concentrations originating in the SoCAB.




Figure 3-6. Wind rose from state 24-hr PM_{10} exceedance days during 18 continuous monitoring years at the Landfill (top right) and Community (bottom right) sites. Wind data at the Community site have been replaced with those from the Reseda site from Year 11 to Year 16 (as discussed in Section 5.1).

Also shown in Figure 3-6 are wind data from the Community site for the 324 state exceedance days during the 18-yr period. On days when 24-hr PM_{10} concentrations exceed the state standard at the

Community site, either (1) wind speeds are relatively low with wind direction predominantly coming from the Los Angeles basin (southeast), or (2) wind speeds are relatively high with wind direction coming from the Landfill (northwest). Regional contributions and landfill activity are thus the main driver of PM₁₀ state exceedances at the Community site. After 18 years of continuous data collection, it is clear that PM₁₀ state exceedances are more common at the Landfill site than they are at the Community site. In addition, differences in wind speed and direction patterns between the two sites on days of measured state exceedances provide insight on the source contributions.

4. Regional Comparisons of PM₁₀

Comparing the PM₁₀ concentrations measured at the Landfill and Community sites with those measured at nearby regional monitoring sites places the locally collected data in a larger, more regional context. The Landfill and Community sites are directly affected by emissions in the SoCAB and the nearby highly trafficked freeway system. The sites chosen for comparison, shown earlier in Figure 1-1, are the closest regulatory sites that conduct routine PM₁₀ monitoring.

Figure 4-1 shows the monthly average PM₁₀ concentrations for the Landfill and Community sites and the three regional locations from January 2008 through November 2025. Note that a 75% data completeness threshold was used when calculating the monthly averages. For example, at least 23 valid daily PM₁₀ values are needed to calculate the monthly average PM₁₀ for a month with 31 days. For the first three years of continuous monitoring, the South Coast AQMD monitor in downtown Los Angeles recorded on average the highest PM₁₀ concentrations among the three regional sites, with exceptions noted in May 2009 and June/July 2010. These exceptions were discussed in the *Third Annual Report of Ambient Air Quality Monitoring at Sunshine Canyon Landfill and Van Gogh Elementary School (June 1, 2009–May 31, 2010)*.⁸ The regional monitor in Burbank followed a month-to-month pattern similar to the Los Angeles pattern, but at a lower average PM₁₀ concentration, until the site was discontinued in summer 2014. The FRM monitor at Santa Clarita on the northern edge of the air basin recorded on average the lowest PM₁₀ concentrations of the regional sites. From 2008 to 2010, Landfill and Community measurements tended to track between the Los Angeles and Santa Clarita data.

The monitoring years since 2011 deviated from this pattern, with the Landfill site usually exhibiting the highest average monthly concentrations from June through September. To help explain this pattern and emphasize the importance of the effect of meteorology on measured pollutant levels, the *Ninth Annual Report of Ambient Air Quality Monitoring at Sunshine Canyon Landfill and Van Gogh Elementary School (November 22, 2007–November 21, 2016)*,⁹ provides meteorological data measured at the Landfill site for the 2008 through 2016 summer seasons. These data demonstrate that measurements at the Landfill site are dominated by wind flow from the south to south-southeast, and thus by regional PM₁₀ concentrations originating in the SoCAB. The dominance of low speed, south-southeasterly winds from June through September between 2011 and 2016 was coupled with PM₁₀ concentrations at the Landfill site that consistently exceeded those of the downtown Los Angeles monitor. The main conclusion drawn from these periods of low-speed southerly winds is that summertime elevations of PM₁₀ concentrations measured at the Landfill site are not solely attributable to landfill activities. A deviation from the pattern occurred in 2017, with the Landfill site exhibiting the highest average monthly concentrations consistently throughout the

⁸ The *Third Annual Report of Ambient Air Quality Monitoring at Sunshine Canyon Landfill and Van Gogh Elementary School (June 1, 2009–May 31, 2010)* was delivered to the Los Angeles City Planning Department in March 2011

⁹ The *Ninth Annual Report of Ambient Air Quality Monitoring at Sunshine Canyon Landfill and Van Gogh Elementary School (November 22, 2007–November 21, 2016)* was delivered to the Los Angeles City Planning Department in April 2017.

year among the sites shown in Figure 4-1. Uncharacteristic monthly concentration spikes occurred at the Landfill site in December 2016, April 2017, and October 2017. In addition, concentrations followed a similar elevated summer season pattern. However, the deviations in monthly average PM₁₀ concentrations from the next highest monitor (South Coast AQMD monitor in downtown Los Angeles) were the largest on record. In 2018, the Landfill site again followed an elevated summer season pattern. In 2019, both the Landfill site and the Community site had an abnormal increase in the monthly average PM₁₀ concentrations during the fall quarter due to wildfire impacts. Year 14 saw an increase in monthly average PM₁₀ concentrations at the Landfill site relative to the previous years of monitoring, with a notable spike of 96 µg/m³ in July 2021, followed by a sharp decrease throughout the remainder of the months and into Year 15. In Year 16, monthly average PM₁₀ concentrations remained high relative to the previous 15 years but were lower relative to the previous three years. Monthly average PM₁₀ concentrations in Year 17 and Year 18 showed a drastic decrease from the previous four years. PM₁₀ concentrations at the Community site remained lower than at the Landfill site and correlated with concentrations recorded in Santa Clarita. [Figure 4-2](#) shows a zoomed-in version of Figure 4-1, highlighting a subset of data from Year 13 to the current year. As shown in Figure 4-2, monthly concentrations at the Landfill site have fallen dramatically from the previous four years and are more similar to regional PM₁₀ sites.

[Figure 4-3](#) shows the rolling annual average PM₁₀ concentrations for the Landfill and Community sites and the three regional locations for Year 1 (2008) through Year 18 (2025). The rolling annual average is calculated from yearly averages over a period of three years. For example, the annual rolling average for Year 18 is calculated based on the individual yearly averages of Years 16, 17, and 18. Due to the nature of this method, the rolling annual average for Year 1 and Year 2 are their yearly averages, respectively. (The yearly averages are calculated based on either hourly or 1-in-6-day FRM measurements.) While Figure 4-1 provides valuable insight on monthly and seasonal variations for each site, a rolling annual average allows for a more concise site-by-site comparison over the entire monitoring period. The Landfill site compares well with the downtown Los Angeles site in Years 5 through 9. However, starting in Year 10, the average concentrations at the two sites deviate from each other, with the Landfill site showing a significant increase. Concentrations at the Landfill site reached a peak in Year 15 and have been trending down over the following three years.

In Years 5 through 7, the PM₁₀ concentrations are consistently higher at the Community site than at the regional monitor in Burbank and the Santa Clarita FRM. However, annual average concentrations are lower at the Community site than at the downtown Los Angeles and Landfill sites for the 18-yr period. Furthermore, the concentrations at the Community site have been steadily decreasing since Year 7, and in Year 11 they fell below the annual average concentrations measured at Santa Clarita until Year 14, when concentrations at the Santa Clarita site rose slightly below those at the Community site. In Year 18, concentrations at the Community site are slightly lower than those at the Santa Clarita site.




Figure 4-1. Monthly average PM₁₀ concentrations for the Landfill, Landfill North (in operation from June 2016 until May 2017), and Community sites, and three regional monitoring sites for 2008–2025. The Santa Clarita site reports integrated 24-hr samples of filters on a one-in-six-day schedule. As of June 30, 2014, the Burbank site is no longer actively reporting PM₁₀ data. The current year (Year 18) time period is shaded in grey.




Figure 4-2. Monthly average PM₁₀ concentrations for the Landfill and Community sites, and three regional monitoring sites from Year 13 through Year 18. The Santa Clarita site reports integrated 24-hr samples of filters on a one-in-six-day schedule. The current year (Year 18) time period is shaded in grey.




Figure 4-3. Rolling 3-yr annual average PM₁₀ concentrations for the Landfill and Community sites, and three regional monitoring sites for Monitoring Year 1–Year 18 (e.g., Monitoring Year 1 refers to Nov. 22, 2007–Nov. 21, 2008). Additional FRM data collected at the Los Angeles site and at the Burbank site are also included.

5. PM₁₀ and BC: Effects of Wind Direction and Work Activity Levels

Both wind direction and landfill work activity levels affect PM₁₀ and BC concentrations measured at the Landfill and Community sites. As described in Sections 3 and 4, winds coming from the south, for example, transport pollutants from densely populated areas of the SoCAB, and have a major effect on local pollutant concentrations. Similarly, landfill contributions to neighborhood-scale PM₁₀ and BC concentrations are expected under northerly wind flow. PM₁₀ and BC concentrations are also expected to vary diurnally, and from day to day, as source strengths increase and decrease with changing activity levels. These activity levels vary with different times of day (e.g., daytime versus nighttime) or between working days and holidays, both regionally and at the local (landfill operations) scale.

The 18-yr data archive is used here with long-term averaging to compare the concentrations of PM₁₀ and BC that characterize the Landfill and Community sites under northerly and southerly wind flows and under differing activity levels (subsections 5.1 to 5.5). Activity levels are binned according to landfill working and non-working days and working and non-working hours. The 18-yr averaged results presented in this report concerning the effect of work activity levels on concentrations of PM₁₀ and BC are, overall, consistent with those presented in Sonoma Technology's third through fourteenth annual reports.

The *Ninth Annual Report of Ambient Air Quality Monitoring at Sunshine Canyon Landfill and Van Gogh Elementary School (November 22, 2007–November 21, 2016)*, provides a comparative analysis of the PM₁₀ and BC levels at the Landfill and Landfill North sites in 2016. Because the Landfill North site operated for only one year, it is not included in subsequent analyses. However, subsection 5.6 of the Ninth Annual Report described the additional comparisons of PM₁₀ and BC concentrations between the Landfill and Landfill North sites by wind direction and landfill work activity levels, and this information is reproduced in Section B.3 of this report (Appendix B).

5.1 General Wind Roses for the Monitoring Sites

Figure 5-1 and Figure 5-2 show 2-yr groups of annual wind roses at the Landfill and Community site from 2007 through 2017 (Years 1 through 10), and individual wind roses for 2018 through 2025 (Years 11 through 18). Wind data from the Community site from Year 11 (i.e., since November 22, 2017) through Year 16 were substituted with data from the nearby Reseda site. While data completeness for wind speed and wind direction at the Community site were above 99% (as depicted in Table 2-1), a database issue¹⁰ prevented the use of the Community wind data since Year

¹⁰ Wind data appears shifted from typical patterns and is currently suspect.

11. The Reseda data were chosen as a surrogate because (1) the Reseda site is operated by South Coast AQMD and follows strict data collection and quality standards; (2) the Reseda site is located just over 4 miles to the south of the Community site, and no topographical barriers exist between it and the Community site; and (3) historical wind patterns at the Reseda site are most representative of the historical patterns of the Community site (i.e., shows the strongest winds from the north and light, variable winds from other directions). For Year 18, it was determined that the quality of wind data was sufficient to use instead of the surrogate.




Figure 5-1. Wind roses for the Landfill site over the 18 years of monitoring data. Wind data for monitoring Years 1 through 16 are shown in 2-yr groups, while the data for Year 17 through Year 18 are displayed as individual wind roses. Only data labeled “valid” or “suspect” are used.




Figure 5-2. Wind roses for the Community Site over the 18 years of monitoring data. Wind data for monitoring Years 1 through 16 are shown in 2-yr groups, while the data for Year 17 through Year 18 are displayed as individual wind roses. Only data labeled “valid” or “suspect” are used. Wind data at the Community site were replaced with those from the Reseda site from Year 11 to Year 16.

Winds at the Landfill site are strongest when they are from the north and north-northwest, and southerly winds are lighter. Community site winds are also strongest from the north-northwest, while winds from all other directions are generally lighter. The wind data show that the winds at the Landfill site are highly directional, and winds at the Community sites are more variable.

Figure 5-3 shows a pollution rose and a pollution differential rose for hourly BC concentration at the Community site. A pollution rose is akin to a bar graph of concentrations associated with wind direction. As shown in Figure 5-3, the lowest hourly BC concentrations at the Community site are

associated with winds from the northwest (as shown in the top graphic). In contrast, the pollution differential rose in the bottom graphic shows that the highest hourly BC concentrations at the Community site (when hourly BC concentrations are higher than those at the Landfill site) are associated with winds from the south and southeast.




Figure 5-3. Top panel: pollution rose of hourly BC concentration at the Community site. Bottom panel: pollution differential rose of excess hourly BC concentration at the Community site (compared to the Landfill site). Data are from December 7, 2007, through November 21, 2025. Data are used only when both BC and wind direction are labeled “valid.”

5.2 Wind Direction Sectors for Categorizing Data

In light of the information about directional winds influencing pollutant concentrations, data for this analysis were selected by using one wind sector to represent the landfill source and areas to the north, and a second wind sector to represent the area from which pollutants travel from the SoCAB.

Figure 5-4 shows the wind sectors representing the landfill source in black for the Landfill site and green for the Community site. The Landfill site's wind sector (greater than or equal to 303 degrees and less than or equal to 360 degrees from true north) is broader than the Community site's (greater than or equal to 325 degrees and less than or equal to 355 degrees from true north). Hourly pollution data corresponding to hourly wind direction data that fall within the boundaries of these sectors are used to compute the pollution metrics for working and non-working days (or hours). The analysis is based only on direction, not on matching times between records at the two sites. The underlying premise is that long-term averages calculated in this manner more accurately represent true average landfill-derived contributions than those calculated from matched hourly records.



Figure 5-4. Aerial image of the Sunshine Canyon Landfill and the surrounding area, showing the wind direction sectors representing the landfill source used to select data for analysis from the Landfill monitor (in black) and the Community monitor (in green).

Figure 5-5 shows the wind sector representing the SoCAB source for both the Landfill and Community sites (greater than or equal to 150 degrees and less than or equal to 210 degrees from true north).




Figure 5-5. Aerial image of the Sunshine Canyon Landfill and the northern portion of the SoCAB, showing the wind direction sector representing the SoCAB source used to select data for analysis to compare with the landfill wind direction sectors depicted in Figure 5-4. The white dot represents the Landfill monitor, and the black dot represents the Community monitor.

5.3 Working and Non-Working Days and Hours for Categorizing Data

After the hourly data have been initially binned by the wind direction sectors described above, hourly PM_{10} and BC concentrations are categorized into the landfill's working and non-working days, and working hours (defined as beginning at 06:00 PST and ending at 17:00 PST) and non-working hours within those days. Working days at the landfill are defined as Monday through Friday, excluding federal holidays. Non-working days are considered Sundays and federal holidays, including New Year's, Memorial, Independence, Labor, Thanksgiving, and Christmas days. Additional non-Sunday holidays when the landfill is closed, but operating, would also be incorrectly binned and thus slightly skew the resulting estimates for that category. Saturdays are categorized "mixed use" at the landfill and thus do not fit easily into either category. The non-Sunday holidays and Saturdays are excluded from the analysis.

5.4 PM_{10} Concentrations

Figure 5-6 provides a visual key for interpreting a notched box-whisker plot. **Figure 5-7** through **Figure 5-10** show notched box-whisker plots that summarize the 18-yr and Year 18 (2025) hourly average PM_{10} concentrations at the Landfill and Community sites for the northerly and southerly wind sectors for working and non-working days and for working and non-working hours within those days. **Figure 5-11** through **Figure 5-14** illustrate median PM_{10} concentrations and 95% confidence intervals for working and non-working days and for working and non-working hours for each wind sector.

A notched box-whisker plot shows the entire distribution of concentrations for each year. In box-whisker plots, each box shows the 25th, 50th (median), and 75th percentiles. The boxes are notched (narrowed) at the median and return to full width at the 95% lower and upper confidence interval values. These plots indicate that we are 95% confident that the median falls within the notch. **Figure 5-15** depicts the hourly average PM_{10} concentrations at the Burbank and Los Angeles regional monitoring sites for working and non-working days and for working and non-working hours within those days in notched box-whisker plots.




Figure 5-6. Instructions for interpreting notched box-whisker plots.

The following general conclusions are based on the results depicted in Figure 5-7 through Figure 5-15.

- During the highest activity levels (working hours on working days):
 - When the wind is from the SoCAB, the Landfill site typically measures slightly higher concentrations of PM_{10} than the Community site. This is true over the 18-yr monitoring period and in Year 18. In Year 18, the differences in PM_{10} concentrations between the Landfill site and the Community site are not statistically significant, as indicated by the overlapping notches in the box-whisker plot. This observation for PM_{10} concentrations in Year 18 is consistent with decreasing concentrations at the Landfill site.
 - At the Community site, the median concentration of PM_{10} when the wind is from the SoCAB is typically about two times higher than when the wind is from the landfill. This is true over the 18-yr period and in Year.
 - When wind is from the landfill, the median PM_{10} concentration at the Community site is approximately one-third of that measured at the landfill, suggesting that although the landfill-derived PM_{10} concentrations are significant, they remain mostly localized to the landfill. This is true when investigating PM_{10} concentrations over the entire 18-yr period and in Year 18.
 - At the Community site, the median concentration of PM_{10} on working days is slightly higher than on non-working days when the wind is from either the landfill or the SoCAB (over the entire 18-yr period). This pattern is similar to median PM_{10} concentrations at the regional sites (Burbank and Los Angeles) for working and non-working days/hours,

suggesting an influence of regional day-of-week and working hours' concentration patterns on the Community site.

- During non-working hours on working days:
 - When the wind is from the SoCAB, the Community site measures higher PM₁₀ concentrations than when wind is from the landfill. This is true when investigating PM₁₀ concentrations over the entire 18-yr period and for Year 18.
 - When the wind is from the landfill over the 18-yr monitoring period, PM₁₀ concentrations are lower at both monitoring sites than when the wind is from the SoCAB, with the Community site characterized by lower concentrations than the Landfill site. This pattern illustrates a localized landfill contribution during times of low activity (nighttime).
 - In Year 18, when the wind is from the direction of the landfill, PM₁₀ concentrations at the Landfill site are slightly lower than the concentrations when the wind is from the SoCAB, suggesting a regional influence. Nevertheless, PM₁₀ concentrations measured at the Community site remain lower than the Landfill site whether wind is from the landfill or SoCAB.
- During the lowest activity levels:
 - The median PM₁₀ concentrations are generally lower on non-working days than on working days, but the extent of the difference is influenced by wind direction. At the Landfill site, the median PM₁₀ concentrations in daytime (working hours) showed a greater proportional decrease on non-working days when wind direction was from the landfill than on non-working days when wind came from the SoCAB. This finding reflects the larger regional PM₁₀ influence of the SoCAB on non-working days.
 - At the Community site, the median PM₁₀ concentrations in daytime (working hours) showed a lesser proportional decrease on non-working days than on working days, and wind direction was less of a factor. Furthermore, lower PM₁₀ concentrations during non-working days versus working days at the Community site coincide with workday/non-workday PM₁₀ concentrations patterns at the regional sites (Burbank and Los Angeles).




Figure 5-7. Notched box whisker plots of 18-yr hourly PM_{10} concentrations for northerly ("From Landfill") wind sectors (as displayed in Figure 5-4) for working days (WD), non-working days (NWD), working hours (WH), and non-working hours (NWH) within those days for the Landfill (red) and Community (blue) monitor sites. Outliers over $125 \mu g/m^3$ are not displayed.




Figure 5-8. Notched box whisker plots of 18-yr hourly PM_{10} concentrations for southerly ("From SoCAB") wind sectors (as displayed in Figure 5-5) for working days (WD), non-working days (NWD), working hours (WH), and non-working hours (NWH) within those days for the Landfill (red) and Community (blue) monitor sites. Outliers over 125 $\mu g/m^3$ are not displayed.




Figure 5-9. Notched box whisker plots of Year 18 (2025) hourly PM_{10} concentrations for northerly (“From Landfill”) wind sectors (as displayed in Figure 5-4) for working days (WD), non-working days (NWD), working hours (WH), and non-working hours (NWH) within those days for the Landfill (red) and Community (blue) monitor sites. Outliers over $125 \mu g/m^3$ are not displayed.




Figure 5-10. Notched box whisker plots of Year 18 (2025) hourly PM_{10} concentrations for southerly ("From SoCAB") wind sectors (as displayed in Figure 5-5) for working days (WD), non-working days (NWD), working hours (WH), and non-working hours (NWH) within those days for the Landfill (red) and Community (blue) monitor sites. Outliers over $125 \mu g/m^3$ are not displayed.




Figure 5-11. Eighteen-yr hourly PM_{10} median concentrations for northerly ("From Landfill") wind sectors (as displayed in Figure 5-4) for working days (WD), non-working days (NWD), working hours (WH), and non-working hours (NWH) within those days for the Landfill (red) and Community (blue) monitor sites. 95% confidence intervals are shown in black.




Figure 5-12. Eighteen-yr hourly PM_{10} median concentrations for northerly ("From SoCAB") wind sectors (as displayed in Figure 5-5) for working days (WD), non-working days (NWD), working hours (WH), and non-working hours (NWH) within those days for the Landfill (red) and Community (blue) monitor sites. 95% confidence intervals are shown in black.




Figure 5-13. Year 18 (2025) hourly PM_{10} median concentrations for northerly (“From Landfill”) wind sectors (as displayed in Figure 5-4) for working days (WD), non-working days (NWD), working hours (WH), and non-working hours (NWH) within those days for the Landfill (red) and Community (blue) monitor sites. 95% confidence intervals are shown in black. Note that confidence intervals are a function of the number of data points; fewer data points lead to a wider interval and less certainty about where the median falls.




Figure 5-14. Year 18 (2025) hourly PM_{10} median concentrations for northerly (“From SoCAB”) wind sectors (as displayed in Figure 5-5) for working days (WD), non-working days (NWD), working hours (WH), and non-working hours (NWH) within those days for the Landfill (red) and Community (blue) monitor sites. 95% confidence intervals are shown in black.




Figure 5-15. Hourly PM_{10} median concentrations for (a) Burbank (2008-2014) and (b) Los Angeles (2008-2025) for working days (WD), non-working days (NWD), working hours (WH), and non-working hours (NWH).

5.5 BC Concentrations

Figure 5-16 through Figure 5-19 summarize the 18-yr and Year 18 (2025) hourly average BC concentrations for the northerly and southerly wind sectors during working and non-working days and during working and non-working hours within those days in a notched box-whisker plot. Similar to the PM_{10} concentration section above, Figure 5-20 through Figure 5-23 illustrate median BC concentrations and 95% confidence intervals for working and non-working days and for working and non-working hours for each wind sector. The following general conclusions are based on the statistical values presented in Figure 5-16 through Figure 5-23.

- During the highest activity levels (working hours on working days):
 - The median concentration of BC measured at both the Landfill and Community sites are higher when the winds are from the SoCAB than when they are from the landfill. This is

- true over the 18-yr monitoring period and in Year 18. However, the differences have decreased in recent years as overall BC concentrations have decreased.
- Over the 18-yr monitoring period and in Year 18, when the wind is from the SoCAB, the Community site measures higher levels of BC concentrations than the Landfill site.
 - When the wind is from the SoCAB, the Community site measures almost four times the median concentration of BC as when the wind is from the landfill over the 18-yr monitoring period. In Year 18, when the wind is from the SoCAB, the Community site measures nearly 4.5 times the median concentration of BC than when the wind is from the landfill.
 - When wind comes from the landfill, the Community BC levels are almost one half of the BC levels measured at the landfill itself over the 18-yr monitoring period. In Year 18, Community BC levels are only 0.079 µg/m³ lower than levels at the Landfill site; this corresponds to lower concentrations at the Landfill site in recent years.
- During the lowest activity levels (non-working days):
 - The median concentrations of BC are generally lower on non-working days than on working days in all categories, but the extent of the difference is influenced by wind direction. Compared to the median BC concentrations during non-working hours on working days, the median BC concentrations during non-working hours on non-working days decreased by a factor of approximately 1.2 (Community site) to 1.3 (Landfill site) when winds were from the landfill. They decreased by about a factor of roughly 1.1 (both sites) when winds were from the SoCAB.
 - On working days, diesel-powered vehicles (trucks and earth-moving equipment) operating at the landfill appear to increase the ambient concentrations of DPM, as determined by the BC measurements from the Landfill site. However, the large metropolitan area of the SoCAB remains the dominant source of DPM. Furthermore, increased BC measurements at the Community site on working days versus non-working days (regardless of wind direction) coincide with known metropolitan area DPM source activity patterns.




Figure 5-16. Notched box whisker plots of 18-yr hourly average BC concentrations for northerly ("From Landfill") wind sectors (as displayed in Figure 5-4) for working days (WD), non-working days (NWD), working hours (WH), and non-working hours (NWH) within those days for the Landfill (red) and Community (blue) monitor sites. Outliers over $2 \mu\text{g}/\text{m}^3$ are not displayed.




Figure 5-17. Notched box whisker plots of 18-yr hourly average BC concentrations for southerly ("From SoCAB") wind sectors (as displayed in Figure 5-5) for working days (WD), non-working days (NWD), working hours (WH), and non-working hours (NWH) within those days for the Landfill (red) and Community (blue) monitor sites. Outliers over $2 \mu\text{g}/\text{m}^3$ are not displayed.




Figure 5-18. Notched box whisker plots of Year 18 (2025) hourly average BC concentrations for northerly ("From Landfill") wind sectors (as displayed in Figure 5-4) for working days (WD), non-working days (NWD), working hours (WH), and non-working hours (NWH) within those days for the Landfill (red) and Community (blue) monitor sites. Outliers over $2 \mu\text{g}/\text{m}^3$ are not displayed.




Figure 5-19. Notched box whisker plots of Year 18 (2025) hourly average BC concentrations for southerly ("From SoCAB") wind sectors (as displayed in Figure 5-5) for working days (WD), non-working days (NWD), working hours (WH), and non-working hours (NWH) within those days for the Landfill (red) and Community (blue) monitor sites. Outliers over $2 \mu\text{g}/\text{m}^3$ are not displayed.




Figure 5-20. Eighteen-yr hourly median BC concentrations for northerly ("From Landfill") wind sectors (as displayed in Figure 5-4) for working days (WD), non-working days (NWD), working hours (WH), and non-working hours (NWH) within those days for the Landfill (red) and Community (blue) monitor sites. 95% confidence intervals are shown in black.




Figure 5-21. Eighteen-yr hourly median BC concentrations for northerly ("From SoCAB") wind sectors (as displayed in Figure 5-5) for working days (WD), non-working days (NWD), working hours (WH), and non-working hours (NWH) within those days for the Landfill (red) and Community (blue) monitor sites. 95% confidence intervals are shown in black.




Figure 5-22. Year 18 (2025) hourly median BC concentrations for northerly ("From Landfill") wind sectors (as displayed in Figure 5-4) for working days (WD), non-working days (NWD), working hours (WH), and non-working hours (NWH) within those days for the Landfill (red) and Community (blue) monitor sites. 95% confidence intervals are shown in black.




Figure 5-23. Year 18 (2025) hourly median BC concentrations for northerly ("From SoCAB") wind sectors (as displayed in Figure 5-5) for working days (WD), non-working days (NWD), working hours (WH), and non-working hours (NWH) within those days for the Landfill (red) and Community (blue) monitor sites. 95% confidence intervals are shown in black.

6. Quantitative Estimates of Landfill Impacts on Ambient Concentrations of PM₁₀ and BC

Quantitative estimates of the impact of landfill operations on neighborhood-scale ambient air quality are required by the original Conditions of Approval (C.10.a) and the nearly identical County Condition 81. Specifically, the conditions require determination of “whether air quality near the Landfill is consistent with the supporting environmental documentation for the City Project (i.e., the City’s Final Supplemental Environmental Impact Report [FSEIR]).” The FSEIR reported the emissions estimates of pollutants likely to result from landfill operations, modeled by the Industrial Source Complex Short Term (ISCST3) regulatory model. Beginning with baseline year data (November 22, 2001–November 21, 2002) and continuing through 2008, no attempt was made to specifically address this requirement, primarily because there is no way to *directly* calculate an appropriate metric. Critically, no pollutant monitoring data were gathered immediately upwind of the landfill to enable accurate estimates of the regional concentrations north of the landfill (and thus unaffected by landfill contributions). While the South Coast AQMD operates a BAM-1020 monitor at the Santa Clarita station, it is configured for PM_{2.5} sampling; these PM_{2.5} data are not directly comparable to the PM₁₀ data provided by the BAM-1020 instruments currently deployed at the Landfill and Community monitoring sites. The Santa Clarita station does employ FRM measurements of PM₁₀ (integrated 24-hr samples on filters) on a one-in-six-day schedule. While 24-hr averaged data from the Landfill PM₁₀ monitor could be compared with the 24-hr integrated data from the FRM samples every sixth day, the low frequency of sampling supports only minimal statistical power for calculating upwind (background) PM₁₀ concentrations. Additionally, the location of the Santa Clarita station relative to the landfill and nearby freeways further complicates the potential for direct application of that data for calculating landfill contributions of PM₁₀. Furthermore, wind direction often changes over 24-hr periods, meaning the 24-hr averages from Santa Clarita are not suitable for apportionment by wind direction.

In Year 9 (2016), the data collected at the Landfill North site provided the opportunity for a more direct measurement of contributions of PM₁₀ and BC from landfill operations. The hourly PM₁₀ and BC concentration data from the Landfill and Landfill North sites, when the measured winds at the Landfill site were from the landfill or the SoCAB, were compared by subtracting the Landfill North site values from the Landfill site values to obtain the differences for each wind direction (i.e., from the landfill or from the SoCAB). A similar analysis was conducted for the Landfill and Community sites to determine whether there was any evidence of landfill contribution to the PM₁₀ and BC concentrations at the Community site. The hourly PM₁₀ and BC concentration data from the Landfill and Community sites when the measured winds at the Community site were from the landfill or from the SoCAB were compared by subtracting the Landfill site values from the Community site values to obtain the

differences by wind direction (i.e., from the landfill or from the SoCAB). Results from the Year 9 difference analysis can be found in Appendix B, but the key takeaway is that the directly measured contribution is more than two times higher than the estimated contribution resulting from the previous data analysis method (which began in the Second Annual Report in 2009¹¹). The previous estimation method is difficult to clearly describe and resulted in a lower contribution estimate of the impact of landfill operations on neighborhood-scale ambient air quality than the direct measurement method did; therefore, this report uses the same approach as the direct measurement method by comparing the difference between the Landfill and Community sites under the two wind sectors (i.e., from the SoCAB and from the landfill).

The following general conclusions are based on the PM₁₀ difference values presented in [Figure 6-1](#) through [Figure 6-4](#).

- The greatest difference in PM₁₀ concentrations between the Landfill and Community sites was observed during periods of highest activity levels (i.e., working hours on working days). Over the 18-yr monitoring period, the median PM₁₀ difference was 25 µg/m³ (mean difference of 41 µg/m³) lower at the Community site when the winds were from the landfill. In Year 18, the median PM₁₀ difference was 26 µg/m³ (mean difference of 41 µg/m³) lower at the Community site when the winds were from the landfill.
- When wind was coming from the landfill, PM₁₀ levels at the Community site were lower than those at the Landfill site for all working categories over the 18 years. Additionally, PM₁₀ concentrations measured at the Community site were lower in each working category when compared to regional PM₁₀ measurements (as shown in [Figure 5-15](#)). A landfill contribution to PM₁₀ concentrations at the Community site was not evident under these wind conditions over the 18 years.
- When the wind was from the SoCAB, the PM₁₀ values at the Community site were slightly lower than the values at the Landfill site in the non-working hour categories. The absolute difference in PM₁₀ values between the Community site and Landfill site is less when winds are from the SoCAB than when winds are from the Landfill, indicating a regional contribution of PM₁₀ from the SoCAB to the Community site. On days in the highest activity level category, the regional contribution of PM₁₀ combined with local landfill contributions, increasing PM₁₀ concentrations at the Landfill site. In Year 18, PM₁₀ levels at the Community site were lower than those at the Landfill site when wind was from SoCAB for all working categories except working hours on non-working days, indicating the combination of increased landfill activity and regional PM₁₀ contribution.

¹¹ Vaughn D.L. and Roberts P.T. (2009) Second annual report of ambient air quality monitoring at Sunshine Canyon Landfill and Van Gogh Elementary School. Prepared for the Planning Department, City of Los Angeles, CA, by Sonoma Technology, Inc., Petaluma, CA, STI-907032-3671-AR, August.




Figure 6-1. Median, mean, and standard deviation of PM₁₀ concentration differences at the Community site versus the Landfill site for northerly ("From Landfill") wind sectors (as displayed in Figure 5-4) for working days (WD), non-working days (NWD), working hours (WH), and non-working hours (NWH) based on the 18-yr monitoring period dataset. Negative values represent lower PM₁₀ concentrations at the Community site.




Figure 6-2. Median, mean, and standard deviation of PM₁₀ concentration differences at the Community site versus the Landfill site for southerly (“From SoCAB”) wind sectors (as displayed in Figure 5-5) for working days (WD), non-working days (NWD), working hours (WH), and non-working hours (NWH) based on the 18-yr monitoring period dataset. Negative values represent lower PM₁₀ concentrations at the Community site.




Figure 6-3. Median, mean, and standard deviation of PM₁₀ concentration differences at the Community site versus the Landfill site for northerly (“From Landfill”) wind sectors (as displayed in Figure 5-4) for working days (WD), non-working days (NWD), working hours (WH), and non-working hours (NWH) based on the Year 18 (2025) dataset. Negative values represent lower PM₁₀ concentrations at the Community site, while positive values represent higher PM₁₀ concentrations.




Figure 6-4. Median, mean, and standard deviation of PM₁₀ concentration differences at the Community site versus the Landfill site for southerly ("From SoCAB") wind sectors (as displayed in Figure 5-5) for working days (WD), non-working days (NWD), working hours (WH), and non-working hours (NWH) based on the Year 18 (2025) dataset. Negative values represent lower PM₁₀ concentrations at the Community site.

The following general conclusions are based on the BC difference values presented in [Figure 6-5](#) through [Figure 6-8](#).

- The greatest BC differences between observations at the Community site and Landfill site were observed during the highest activity levels (working hours on working days). Over the 18-yr monitoring period, the median BC difference was 0.170 µg/m³ (mean difference of 0.261 µg/m³) lower at the Community site when the winds were from the landfill. Median BC concentrations during working hours on non-working days were equal between the Community and Landfill sites when winds were from the landfill. During the highest activity levels in Year 18, the median BC difference was 0.079 µg/m³ (mean difference of 0.090 µg/m³) lower at the Community site when the winds were from the landfill, indicating no evident landfill contribution to BC levels at the Community site.

- BC concentrations were higher at the Community site than at the Landfill site when the wind was from the SoCAB in the working hour categories, but they were slightly lower during non-working hour categories. This suggests increased regional BC concentrations contributing to BC levels at the Community site. In Year 18, BC concentrations were higher in all categories at the Community site than at the Landfill site when winds were from the SoCAB except for the working day, non-working hour category.




Figure 6-5. Median, mean, and standard deviation of BC concentration differences at the Community site versus the Landfill site for northerly (“From Landfill”) wind sectors (as displayed in Figure 5-4) for working days (WD), non-working days (NWD), working hours (WH), and non-working hours (NWH) based on the 18-yr monitoring period dataset. Negative values represent lower BC concentrations at the Community site, while positive values represent higher BC concentrations.




Figure 6-6. Median, mean, and standard deviation of BC concentration differences at the Community site versus the Landfill site for southerly ("From SoCAB") wind sectors (as displayed in Figure 5-5) for working days (WD), non-working days (NWD), working hours (WH), and non-working hours (NWH) based on the 18-yr monitoring period dataset. Negative values represent lower BC concentrations at the Community site, while positive values represent higher BC concentrations.



Figure 6-7. Median, mean, and standard deviation of BC concentration differences at the Community site versus the Landfill site for northerly (“From Landfill”) wind sectors (as displayed in Figure 5-4) for working days (WD), non-working days (NWD), working hours (WH), and non-working hours (NWH) based on the Year 18 (2025) dataset. Negative values represent lower BC concentrations at the Community site, while positive values represent higher BC concentrations.




Figure 6-8. Median, mean, and standard deviation of BC concentration differences at the Community site versus the Landfill site for southerly (“From SoCAB”) wind sectors (as displayed in Figure 5-5) for working days (WD), non-working days (NWD), working hours (WH), and non-working hours (NWH) based on the Year 18 (2025) dataset. Negative values represent lower BC concentrations at the Community site, while positive values represent higher BC concentrations.

7. Routine Field Operations

Field operations include regular visits to both monitoring sites. During the first four years of the study, these visits were scheduled at two-week intervals. We changed this to monthly intervals because experience demonstrated that monthly visits suffice to meet the routine maintenance operations associated with the BAM and the Aethalometer. This protocol is in keeping with the maintenance schedule recommended by Met One (manufacturer of the BAM) and Magee Scientific (manufacturer of the Aethalometer). This protocol is accompanied by daily review of data that allows problems to be detected quickly. Many times, the detected problems can be addressed remotely via cellular connection to the site instruments. Occasionally, non-scheduled on-site visits by a Spheros technician are required and occur as soon as reasonably possible.

Each quarterly report contains tables with the dates and times of each site visit and a summary of activities that took place. Consult these reports for a summary of field activities that occurred in Years 1 through 18. [Table 7-1](#) and [Table 7-2](#) summarize all visits during Year 18 for the monitoring sites.

Table 7-1. Sunshine Canyon Landfill monitoring site visits and field maintenance and operations in Year 18.

Date of Site Visit	Description of Work
12/26/2024	Meteorological instrument performance audit BAM flow check
1/3/2025	Replaced anemometer on meteorological tower Changed tape on BAM
3/24/2025	Changed tape on BAM
4/1/2025	Replaced monitor
5/20/2025	Checked tape on BAM
5/29/2025	Changed tape on BAM Checked tape on Aethalometer Performed flow checks on BAM and Aethalometer Removed mice from shelter and sealed trailer to prevent more mice from entering
6/5/2025	Replaced tape on BAM
8/20/2025	Replaced tape on BAM Maintenance on Aethalometer
8/26/2025	Maintenance on Aethalometer
9/16/2025	Checked flow on Aethalometer and BAM
10/2/2025	Trailer maintenance

Date of Site Visit	Description of Work
11/4/2025	Checked flow on Aethalometer and BAM Cleaned nozzle, vane, and roller on BAM Replaced tape on BAM

Table 7-2. Community monitoring site visits and field maintenance and operations in Year 18.

Date of Site Visit	Description of Work
1/3/2025	Rebooted router
3/24/2025	Changed tape on BAM
5/20/2025	Checked tape on BAM
5/29/2025	Changed tape on BAM Checked tape on Aethalometer Performed flow checks on BAM and Aethalometer
6/5/2025	Restarted BAM
6/17/2025	Investigated BAM sampling issues Restarted BAM
6/18/2025	Investigated BAM sampling issues Replaced battery on BAM Restarted BAM
6/24/2025	Investigated Aethalometer flow sampling issues Investigated BAM sampling issues Replaced battery on BAM Restarted BAM
6/30/2025	Updated data logger software
7/2/2025	Replaced Aethalometer
8/1/2025	Site maintenance
8/30/2025	Replaced tape on BAM
9/10/2025	Aethalometer maintenance
9/16/2025	Checked flow on Aethalometer and BAM
9/24/2025	Replace Aethalometer
10/2/2025	Tailer maintenance
10/13/2025	Investigated telecommunication issues
10/15/2025	Changed internet router

Date of Site Visit	Description of Work
11/4/2025	Checked flow on Aethalometer and BAM Replaced tape on BAM
11/18/2025	Anemometer calibration and maintenance

Appendix A. Regional Concentrations of Black Carbon

This Appendix contains an analysis of regional concentrations of Black Carbon (BC) from the Multiple Air Toxics Exposure Study V (MATES V).

Concentrations of BC by month and time of day, and a differential between the Landfill and Community sites, are shown in [Figure A-1](#). These data are from the time period of the MATES V study in May 2018 – April 2019. Concentrations of BC are highest in the summer, with a maximum median concentration occurring in September at the Landfill site and in August at the Community site, although the differences between the two months are not statistically different. Concentrations of BC are highest in the early morning hours of 6:00 a.m. to 8:00 a.m. LST (Figure A-1, bottom). The slight diurnal rise in the differential in the early morning hours at 6:00 a.m. LST to 7:00 a.m. LST may indicate a pattern of slightly higher local concentrations at the Community site in the early morning hours.




Figure A-1. Concentrations of BC at the Landfill site (a, d), Community site (b, c), and difference between concentrations at the Community site and Landfill site (c, f) by month (top three figures) and time of day (bottom three figures) for the time period of the MATES V study (April 2018–May 2019). Differentials are shown on the far right; concentrations below zero indicate that concentrations were higher at the Landfill site than at the Community site. Note that outliers above 5 $\mu\text{g}/\text{m}^3$ in figures a, b, d, and e, and outliers above 2.5 $\mu\text{g}/\text{m}^3$ and below -2.5 $\mu\text{g}/\text{m}^3$ in figures c and f, are excluded.

To place the data in a regional context, Landfill and Community BC concentrations during the MATES V period (May 2018 – April 2019) are shown in comparison to MATES V BC measurements that were made at the Burbank, Los Angeles, Pico Rivera, and Huntington Park sites. [Figure A-2](#) shows a comparison of concentrations for the days and hours when each of the sites had valid BC data available during this time period. Concentrations at the Landfill site (SBS) and Community site (VGS) are shown in yellow, while other nearby Los Angeles sites are shown in blue. Median concentrations at the Landfill and Community sites are lower than those measured at the other four sites during the same time period. Moreover, 75th percentile (top of the box) and upper percentile concentrations (indicated by error bars) are also lower at the Landfill and Community sites than at other sites in the Los Angeles Basin.




Figure A-2. A comparison of regional BC concentrations from May 2018 through April 2019 at landfill sites (yellow) and MATES V monitoring stations (blue). In MATES V documentation, Los Angeles is referred to as “Central L.A.” and Burbank is referred to as “Burbank Area”. Note that outliers above $10 \mu\text{g}/\text{m}^3$ are excluded.

Appendix B. Additional Analyses

This appendix contains discussions of the temporal variability in BC, PM₁₀, and wind direction (Section B.1), and of the effects of wind direction and work activity on BC and PM₁₀ (Section B.2). Section B.3 provides information about the Landfill North site, as previously reported in Section 5.6 of the Ninth Annual Report.

B.1 Temporal Variability in BC, PM₁₀, and Wind Direction

As shown in [Figure B-1](#), the diurnal profiles of PM₁₀ and BC are characterized by a morning peak in concentrations at both monitoring locations. The peak in BC occurs between 6:00 a.m. and 8:00 a.m., while the peak in PM₁₀ is broader, occurring between 6:00 a.m. and 10:00 a.m. Since Year 13, a secondary peak in PM₁₀ has been occurring between 5:00 p.m. and 6:00 p.m. at the Landfill site. However, this secondary peak is diminished in Year 18. The diurnal profiles of PM₁₀ and BC at both sites have been generally decreasing over the past five years of monitoring. Overall, the mean hourly concentrations of both PM₁₀ and BC are lower at the Community site than at the Landfill site.

As shown in the box-whisker plots ([Figure B-2](#)), median concentrations of PM₁₀ and BC are higher during the warm season (approximately May through September) at both the Community and the Landfill sites. The median PM₁₀ concentrations peak in June at both sites. Median BC concentrations are roughly equal in August and September at the Landfill site and are roughly equal in June through September at the Community site.

[Figures B-3](#) through [B-5](#) show seasonal wind roses of hourly wind data collected at the Landfill and Community sites. At the Landfill site, winds are predominantly from the northerly and southerly directions during all seasons, with a larger proportion of winds from the north during the winter and from the south during the summer ([Figures B-3](#) and [B-4](#)). At the Landfill North site, the prevailing winds are northwesterly in the winter, southerly in the spring and summer, and a mix of northwesterly and southerly in the fall (not shown). The prevailing wind direction at the Community site varies during all seasons ([Figure B-5](#)).




Figure B-1. Mean PM_{10} (a, b) and BC (c, d) concentrations by hour for the 18 monitoring years at the Landfill (a, c) and Community (b, d) sites. *Note: the upper limit on the y-axis of plot c is half that of plot a.*




Figure B-2. Distribution of daily mean PM₁₀ and BC concentrations by month across all 18 monitor years (2007–2025) at the Landfill (a, b) and Community (c, d) sites. Note: BC outliers greater than 3 µg/m³ and PM₁₀ outliers greater than 200 µg/m³ are excluded.




Figure B-3. Percent of time during which winds at the Landfill (a) and Community (b) monitoring sites originated from each wind direction sector (South Coast Air Basin, Landfill, Other) during each month across all 18 years (2007–2025).




Figure B-4. Seasonal wind roses based on hourly data collected at the Landfill site from 2007-2025.




Figure B-5. Seasonal wind roses based on hourly data collected at the Community site from 2007-2025. Data from Year 11 to Year 16 are replaced with wind data from the Reseda site.

B.2 BC and PM₁₀: Effects of Wind Direction and Work Activity Levels

As shown in **Figure B-6**, concentrations of BC and PM₁₀ are higher on weekdays than weekends. Higher concentrations are consistent with greater activity at the landfill during the week, as well as with more vehicles on the roads throughout the SoCAB. Concentrations of BC and PM₁₀ are higher on Saturdays than Sundays at the Landfill site. Activity occurs at the landfill on some Saturdays, but not on Sundays.




Figure B-6. Hourly BC (left) and PM₁₀ (right) concentrations at the Landfill (SBS), Landfill North (SBSU), and Community (VGS) monitoring sites on weekdays (blue), Saturdays (pink), and Sundays (green) from November 22, 2007, through November 21, 2025. Note that this plot includes the Landfill North site, which closed in May 2017. Note that BC data greater than 5 µg/m³ and PM₁₀ data greater than 200 µg/m³ are excluded.

As shown in **Figure B-7**, concentrations of BC and PM₁₀ are several times greater when winds come from the south than from the north. In addition, concentrations are typically similar between the Landfill and Community sites when winds are from the SoCAB direction. Concentrations are greater at the Landfill site than the Community site when winds are from the north.




Figure B-7. PM₁₀ (left) and BC (right) concentrations at the Landfill (dark blue) and Community (light blue) sites from November 22, 2007, through November 21, 2025, when winds originated from the Landfill versus when they originated from the SoCAB. Results are based on hourly data points where both sites experienced winds from the same sector. Note that BC data greater than 5 µg/m³ and PM₁₀ data greater than 200 µg/m³ are excluded.

B.3 PM₁₀ and BC: Landfill vs. Landfill North and Community Sites

The data collected at the new Landfill North site in the ninth and tenth monitoring years provided an opportunity to further investigate and characterize the impacts of wind direction and landfill work activity levels on the measured PM₁₀ and BC concentrations at the Landfill site. The hourly PM₁₀ and BC concentration data from the Landfill and Landfill North sites when the measured winds at the Landfill site were from the landfill and from the SoCAB were compared by subtracting the Landfill North site values from the Landfill site values to obtain the differences. The results for PM₁₀ and BC are shown in [Figures B-8](#) and [B-9](#), respectively.


The following general conclusions are based on the median PM₁₀ difference values presented in [Figure B-8](#).

- The greatest difference between the Landfill and Landfill North sites was observed during the periods of highest activity (i.e., working hours on working days, panel a). The PM₁₀ differences were 22 and -26 µg/m³ when the winds were from the landfill and from the SoCAB respectively, suggesting a consistent localized PM₁₀ contribution of 20 to 25 µg/m³ from the landfill to the Landfill site downwind.


- When the wind was from the landfill, the PM₁₀ values were higher at the Landfill site (downwind) than the values at the Landfill North site (upwind) in all working categories, indicating a localized contribution of PM₁₀ from the landfill to the Landfill site.
- When the wind was from the SoCAB, the PM₁₀ values were higher at the Landfill North site (downwind) than the values at the Landfill site (upwind) in all but the non-working hours on non-working days' category, indicating a localized contribution of PM₁₀ from the landfill to the Landfill North site. The median difference for the non-working hours on non-working days' category was zero with a negative mean of -0.2 µg/m³.

The following general conclusions are based on the median BC difference values presented in Figure B-9:


- During the highest activity levels (working hours on working days, panel a), the greatest BC differences were observed. The BC differences were 0.1 µg/m³ when the winds were from the landfill and -0.3 µg/m³ from the SoCAB, suggesting a localized BC contribution from activities at the landfill to the Landfill site downwind. This is the only category where the downwind site showed higher BC concentrations than the upwind site.
- During the time periods of the other working categories, although the median concentrations were slightly higher at the upwind monitor, the BC levels between the two sites were mostly very similar regardless of wind direction. The only exception is that the Landfill site measured notably higher BC when the wind came from SoCAB during non-working hours on non-working days (panel d).
- **Figure B-10** provides an illustration of landfill impact on PM₁₀ and BC concentrations at the downwind site when wind is from either the landfill or the SoCAB as measured at the Landfill site during working hours on working days.




(a) Working hours on working days



(b) Non-working hours on working days




(c) Working hours on non-working days




(d) Non-working hours on non-working days


Figure B-8. Notched box plots of the differences in PM₁₀ concentrations between the Landfill North and the Landfill sites (Landfill site values–Landfill North site values) for northerly and southerly wind sectors for working and non-working days and for working and non-working hours within those days. Outliers over $\pm 100 \mu\text{g}/\text{m}^3$ are not displayed.




(a) Working hours on working days



(b) Non-working hours on working days



(c) Working hours on non-working days



(d) Non-working hours on non-working days

Figure B-9. Notched box plots of the differences in BC concentrations between the Landfill North and the Landfill sites (Landfill site values–Landfill North site values) for northerly and southerly wind sectors for working and non-working days and for working and non-working hours within those days. Outliers over $\pm 1 \mu\text{g}/\text{m}^3$ are not displayed.




Figure B-10. Map depicting the localized impact of the landfill on PM₁₀ and BC concentrations when the wind is from the landfill or the SoCAB as measured at the Landfill site during working hours on working days.

Appendix C. Comparison of Ambient Air Toxics Concentrations to the Final Supplemental Environmental Impact Report

The city's Final Supplemental Environmental Impact Report (FSEIR) reported emissions estimates of pollutants likely to result from landfill operations, modeled by the Industrial Source Complex Short Term (ISCST3) regulatory model. The reported pollutants included a number of hazardous air pollutants (HAPs) but did not include criteria pollutants such as PM_{2.5} or DPM. One year of HAPs measurements were collected at the Landfill and Community sites on a one-in-six-day EPA sampling schedule from July 2016 through June 2017. Target HAPs included key air toxics in the MATES IV protocol, such as benzene, tetrachloroethene, 1,3-butadiene, carbon tetrachloride, dichloromethane, ethylbenzene, xylenes, toluene, and trichloroethene, as well as tracers of landfill emissions such as chlorobenzene, dichlorobenzenes, and vinyl chloride.¹

Table C-1 shows the average concentrations in parts per billion (ppb) measured at both sites during the HAPs measurement campaign. It also shows the average difference between the two sites and the percent difference; in both cases, a negative number indicates the Community site had a higher average concentration than the Landfill site. The FSEIR annual increment shows the modeled annual increment of each pollutant at the "Maximum exposed individual" residence in units of ppb. In almost all cases, the concentration differences and increments are small in absolute terms, with concentrations at or below a few parts per trillion (ppt). The increment is typically much less than 10% of the Community site average. Given the method detection limit of the sampling and analytical methodology of ~6-10 ppt, differences of a few ppt are too small to reliably detect because the sensitivity of the instrument is a few times larger than the value we are trying to detect. The overall concentrations of HAPs in the Sunshine Canyon area are lower than those in most other places in the Los Angeles basin.

¹ McCarthy M.C., O'Brien T.E., Vaughn D.L., Penfold B.M., and Hafner H.R. (2017) Sunshine Canyon VOC and carbonyl monitoring report. Final report prepared for the City of Los Angeles Planning Department, Los Angeles, CA, and the Los Angeles County Department of Regional Planning, Los Angeles, CA, by Sonoma Technology, Inc., Petaluma, CA, STI-916007-6823-FR, November.

Table C-1. Summary statistics for average concentrations (ppb) and differences between the two monitoring sites for HAPs with at least one measurement above detection at the Landfill and Community sites from July 2016 through June 2017. Negative differences indicate values at the Community site are higher than those at the Landfill site. N/A indicates that the HAP is not modeled or reported in the FSEIR.

Parameter	Landfill Site Avg. (ppb)	Community Site Avg. (ppb)	Average Difference (ppb)	% Difference	FSEIR Annual Increment (ppb)
1,2-Dibromoethane	0.009	0.005	0.003	47.2	0.0000013
1,2-Dichlorobenzene	0.009	0.008	0.001	13.6	0.0018 ^a
1,2-Dichloropropane	0.005	0.005	0	-1.9	N/A
1,3-Dichlorobenzene	0.007	0.006	0.001	11.6	0.0018 ^a
1,4-Dichlorobenzene	0.013	0.014	0	-3.3	0.0018 ^a
Benzene	0.171	0.154	0.017	10.2	0.0019
Benzyl chloride	0.005	0.005	0	-1	0.00029
Carbon tetrachloride	0.104	0.106	-0.003	-2.4	0.000013
Chlorobenzene	0.006	0.006	0	-2.2	0.00060
Chloroform	0.017	0.022	-0.006	-28.4	0.0001
Dichloromethane	0.059	0.057	0.002	3.5	0.0043
Ethylbenzene	0.051	0.06	-0.009	-15.4	N/A
Formaldehyde	2.166	2.087	0.079	3.7	N/A
m,p-Xylenes	0.17	0.195	-0.026	-14	0.033 ^b
o-Xylene	0.051	0.06	-0.009	-16.5	0.033 ^b
Styrene	0.028	0.025	0.003	13.1	N/A
Tetrachloroethene	0.019	0.012	0.007	44.4	0.0022
Toluene	0.295	0.285	0.01	3.5	0.042
Trichloroethene	0.007	0.006	0.001	16.1	0.000066

Notes:

^a FSEIR reported only dichlorobenzene without specifying the isomer. Here, we report the sum of dichlorobenzene isomers as shown in FSEIR, but it may be more appropriate to divide the reported 0.0018 ppb increment by three to indicate the individual isomer contributions if they occur in equal portions.

^b FSEIR reported the sum of o, m, and p-xylene isomers as 0.033 ppb. A better estimate is the contribution is 2/3 m- and p-xylene, and 1/3 o-xylene.